

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 9/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/37181</b> <b>(43) International Publication Date:</b> 27 August 1998 (27.08.98)															
<b>(21) International Application Number:</b> PCT/US98/03404 <b>(22) International Filing Date:</b> 20 February 1998 (20.02.98)  <b>(30) Priority Data:</b> <table border="0"><tr><td>60/038,750</td><td>20 February 1997 (20.02.97)</td><td>US</td></tr><tr><td>60/047,151</td><td>20 May 1997 (20.05.97)</td><td>US</td></tr><tr><td>60/054,549</td><td>1 August 1997 (01.08.97)</td><td>US</td></tr><tr><td>60/055,762</td><td>14 August 1997 (14.08.97)</td><td>US</td></tr><tr><td>60/064,322</td><td>30 October 1997 (30.10.97)</td><td>US</td></tr></table> <b>(71) Applicant (for all designated States except US):</b> WHITEHEAD INSTITUTE FOR BIOMEDICAL RESEARCH [US/US]; Nine Cambridge Center, Cambridge, MA 02142 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> COUNTER, Christopher, M. [CA/US]; Apartment 5R, 310 Prospect Street, Cambridge, MA 02139 (US). MEYERSON, Matthew [US/US]; 11 Aqueduct Road, Wayland, MA 01778 (US). WEINBERG, Robert, A. [US/US]; 25 Copley Street, Brookline, MA 02146 (US).  <b>(74) Agents:</b> GRANAHAN, Patricia et al.; Hamilton, Brook, Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA 02173 (US).		60/038,750	20 February 1997 (20.02.97)	US	60/047,151	20 May 1997 (20.05.97)	US	60/054,549	1 August 1997 (01.08.97)	US	60/055,762	14 August 1997 (14.08.97)	US	60/064,322	30 October 1997 (30.10.97)	US	<b>(81) Designated States:</b> CA, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
60/038,750	20 February 1997 (20.02.97)	US															
60/047,151	20 May 1997 (20.05.97)	US															
60/054,549	1 August 1997 (01.08.97)	US															
60/055,762	14 August 1997 (14.08.97)	US															
60/064,322	30 October 1997 (30.10.97)	US															
<b>(54) Title:</b> TELOMERASE CATALYTIC SUBUNIT GENE AND ENCODED PROTEIN  <b>(57) Abstract</b> <p>Isolated DNA encoding the catalytic subunit of a eukaryotic telomerase holoenzyme, such as the catalytic subunit of a yeast or human telomerase holoenzyme; the RNA transcript, which is expressed in primary human tumors, cancer cell lines and telomerase-positive tissues; and the encoded catalytic subunit protein. Methods of assessing cells for malignancy or an increased likelihood of progression to malignancy and methods of diagnosing or aiding in the diagnosis of development of malignancy in an individual are also described.</p>																	

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

TELOMERASE CATALYTIC SUBUNIT GENE AND  
ENCODED PROTEIN

FUNDING

Work described herein was funded, in whole or in part,  
5 by Grant No. R35CA39826 and Grant No. OIG CA 39826 from the  
National Cancer Institute. The U.S. Government has certain  
rights in the invention.

RELATED APPLICATIONS

This application claims the benefit of U.S.  
10 Provisional Application No. 60/038,750, filed February 20,  
1997, entitled "Identification of a Protein Subunit of the  
Yeast Telomerase Holoenzyme" by Christopher M. Counter,  
Matthew Meyerson and Robert A. Weinberg; of U.S.  
Provisional Application No. 60/047,151, filed May 20, 1997,  
15 entitled "Human Telomerase Catalytic Subunit" by  
Christopher M. Counter and Matthew Meyerson; of U.S.  
Provisional Application No. 60/054,549, filed August 1,  
1997, entitled "hEST2, the Putative Human Telomerase  
Catalytic Subunit Gene Is Up-Regulated in Tumor Cells and  
20 During Immortalization" by Matthew Meyerson, Christopher M.  
Counter and Robert A. Weinberg; of U.S. Provisional

-2-

Application No. 60/055,762, filed August 14, 1997, entitled "Human Telomerase Catalytic Subunit Gene and Uses Therefor" by Matthew Meyerson, Christopher M. Counter and Robert A. Weinberg; and of U.S. Provisional Application No. 5 60/064,322, filed October 30, 1997, entitled "Human Telomerase Catalytic Subunit Gene and Uses Therefor", by Matthew Meyerson, Christopher M. Counter and Robert A. Weinberg. The entire teachings of these five referenced applications are expressly incorporated herein by 10 reference.

#### BACKGROUND OF THE INVENTION

In most eukaryotes, the telomere ends of linear chromosomes are replicated by the ribonucleoprotein enzyme telomerase (Blackburn, E.H. *Annu. Rev. Biochem.* 53, 163-94 15 (1984); Zakian, V.A., *Science* 270, 1601-7 (1995); Greider, C.W. & Blackburn, E.H., *Cell* 43, 405-13 (1985)). The RNA subunit of this enzyme has now been described in multiple species (Blasco, M.A. et al., *Science* 269, 1267-70 (1995); Feng, J., et al., *Science* 269, 1236-41 (1995); Greider, 20 C.W. & Blackburn, E.H., *Nature* 337, 331-7 (1989); Lingner, J. et al., *Genes Dev.* 8, 1984-98 (1994); McEachern, M.J. & Blackburn, E.H., *Nature* 376, 403-9 (1995); Melek, M. et al., *Mol. Cell Biol.* 14, 7827-38 (1994); Shippen-Lentz, D. & Blackburn, E.H., *Science* 247, 546-52 (1990); Singer, M.S. 25 & Gottschling, D.E., *Science* 266, 404-9 (1994)). Candidate protein subunits have been isolated in the ciliate *Tetrahymena* (Collins, K. et al., *Cell* 81, 677-86 (1995) but not, however, in any genetically tractable organism.

Telomerase is repressed in normal human somatic cells 30 but is re-activated during tumor progression. This re-

-3-

activation is not reflected by changes in the levels of previously cloned genes encoding telomerase subunits.

#### SUMMARY OF THE INVENTION

Described herein are genes which are required for  
5 telomerase enzymatic activity and the respective encoded  
messenger RNAs (transcripts) and proteins. As described  
below, the present invention relates to DNA which is  
involved in telomere length regulation in eukaryotes and  
proteins which are physically associated with the  
10 respective active telomerase enzyme and, thus, are each a  
component of the respective telomerase holoenzyme. In one  
embodiment, the gene, RNA transcript and encoded protein  
are a yeast gene and its encoded RNA transcript and  
protein. In a second embodiment, the gene, RNA transcript  
15 and encoded protein are human. Also described herein is a  
human cDNA which encodes the human telomerase catalytic  
protein subunit.

The yeast telomerase gene is present on chromosome XII  
and its disruption alters telomere maintenance, as  
20 demonstrated by the decrease in telomere length in  
transposon-insertion mutants. The sequence of the gene,  
referred to herein as *EST2* (SEQ ID NO.: 1), the deduced  
amino acid sequence of the encoded *Est2* protein, also  
referred to as *Est2p*, (SEQ ID NO.: 2) and the *hEST2* RNA  
25 transcript (SEQ ID NO.: 36) are provided. The DNA has been  
shown to be essential for telomerase activity in yeast and  
the encoded protein has been shown, by the methods  
described herein, to be physically associated with  
telomerase and a constituent of active telomerase complex  
30 in yeast. The yeast telomerase DNA *EST2* is a subject of

-4-

the present invention; this includes isolated DNA comprising DNA selected from the following: DNA of SEQ ID NO.: 1; DNA which is the complement of SEQ ID NO.: 1, DNA which hybridizes to DNA of SEQ ID NO.: 1 or a complement thereof; DNA which localizes to yeast chromosome XII; and DNA which encodes the amino acid sequence of SEQ ID NO.: 2. Isolated yeast Est2p is also the subject of this invention. This includes isolated Est2p of SEQ ID NO.: 2 and Est 2p encoded by EST2 DNA as defined herein.

10       The gene described herein is useful to identify genes encoding telomerase proteins in other eukaryotes, particularly in vertebrates, including mammals and especially humans. All or a portion of the gene described can be used. The encoded portion can be used to produce antibodies (monoclonal or polyclonal) which bind the Est2 protein and can, in turn, be used to identify corresponding proteins (proteins physically associated with telomerase enzymatic activity) in other eukaryotes.

Also described herein is a human cDNA, which was originally named *hEST2* and has been renamed *hTERT*, that encodes the human telomerase catalytic protein subunit, the encoded *hEST2* RNA and the encoded *hEST2* protein. The human DNA described herein is a human homologue of yeast and ciliate genes which encode telomerase catalytic subunits; it shares significant sequence similarity with the telomerase catalytic subunit genes, yeast *EST2* and *Euplotes* p123. *hEST2* RNA expression reflects telomerase activity. The RNA transcript is expressed in primary human tumors, cancer cell lines and telomerase-positive tissues, but is undetectable in telomerase-negative cell lines and differentiated telomerase-negative tissues. *hEST2* message is absent in pre-crisis, telomerase-negative transformed cells, but is readily detectable in post-crisis,

-5-

telomerase-positive immortalized cells. Taken together, these observations are evidence that the induction of *hEST2* mRNA expression is required for the telomerase activation that occurs during cellular immortalization and tumor  
5 progression.

The encoded human protein comprises six of the seven conserved sequence motifs which define the polymerase domains of members of the reverse transcriptase family and also includes the invariant aspartic acid residues required  
10 for telomerase enzymatic activity. Although *hEST2* protein comprises such motifs which define polymerase domains, beyond these it shows no sequence similarity with reverse transcriptases. It is more closely related to the telomerase catalytic subunits of yeast and ciliates than to  
15 other reverse transcriptases. In its domains that lie N-terminal to the polymerase domain, *hEST2* shows clear relatedness to both p123 and Est2p. Many of the sequence identities in the N-termini of the three proteins are in a region just before motif 1. These sequences do not appear  
20 in reverse transcriptases or in other proteins and, thus, appear to be unique to telomerases. For example, these three include a unique motif, referred to as the telomerase motif; in *hEST2* this motif extends from amino acid residue 556 to amino acid residue 565 of SEQ ID NO.: 3, with  
25 absolute invariant sequence extending from amino acid residue 560 to amino acid residue 565 of SEQ ID NO.: 3. Further, within the *hEST2* domain that shares sequence similarity with reverse transcriptases, it is clear that *hEST2* is more closely related to telomerase reverse  
30 transcriptases than to non-telomerase reverse transcriptases. The three telomerase catalytic subunits (*hEST2*, yeast *EST2* and *Euplotes* p123) form a subgroup within the reverse transcriptase family.

-6-

The present invention relates to isolated DNA which encodes an RNA transcript which is expressed in primary human tumors, cancer cell line and telomerase-positive tissue. In one embodiment, the isolated DNA comprises

5 *hEST2* DNA such as DNA comprising SEQ ID NO.: 35 (DNA comprising the nucleotide sequence of SEQ ID NO.: 35). Isolated DNA of the present invention encodes an RNA transcript which is not detectable in telomerase-negative cell lines or in differentiated telomerase-negative

10 tissues. Further, isolated DNA of the present invention encodes an RNA transcript which is not detectable in pre-crisis, telomerase-negative transformed cells and is detectable in post-crisis, telomerase-positive immortalized cells. In a particular embodiment, the DNA comprises *hEST2*

15 cDNA, such as SEQ ID NO.: 35 or DNA which hybridizes thereto or to a complement of *hEST2* cDNA (such as a complement of SEQ ID NO.: 35) under conditions of high stringency. A further subject of the present invention is an isolated RNA transcript encoded by isolated DNA of the

20 present invention.

A method of altering telomerase function and, thus, of altering telomere shortening is also described. In the method, expression and/or function of the telomerase-associated protein is altered (enhanced or reduced),

25 resulting in altered (enhanced or reduced) cell lifespan. Expression of the telomerase-associated protein is enhanced, for example, by introducing DNA encoding the protein into cells. Alternatively, telomerase-associated protein is introduced into cells. Expression of the

30 telomerase-associated protein is reduced, for example, by introducing into cells an agent which inhibits production or function of the protein, an agent which destroys the expressed protein or a dominant negative form of the



-7-

protein. Therefore, the present invention also provides a method of increasing the lifespan of a cell or, alternatively, of reducing the lifespan of a cell, such as that of cancer cells or transformed cells. In the method  
5 of increasing lifespan of cells, telomerase activity is maintained or increased within the cells, such that telomeres are maintained and as the cells age, telomere shortening does not occur or occurs to a lesser extent than would otherwise be the case (if telomerase activity were  
10 not maintained or increased). Reduction of lifespan of cells, such as tumor cells, is accomplished by introducing into the cells an inhibitor of the telomerase protein subunit described herein.

A method of assessing cells or aiding in the  
15 assessment of cells for malignancy or an increased likelihood of progression to malignancy is also the subject of this invention. In the method, cells to be assessed are obtained from an individual (e.g., a human) in need of such an assessment. The cells are processed in such a manner  
20 that DNA, RNA or both in the cells are rendered available for annealing or hybridization with complementary polynucleotides or oligonucleotides (DNA or RNA), such as probes or primers, thereby producing processed cells. The processed cells are combined with DNA or RNA required for  
25 telomerase enzymatic activity (DNA or RNA encoding a protein required for telomerase enzymatic activity), a complement of the required DNA or RNA or a characteristic portion or fragment of the DNA or RNA, such as the telomerase motif or all or a portion of the 5' end of *hEST2*  
30 which is not shared with the yeast or *Euplotes* gene. If hybridization occurs, it is indicative of the presence of telomerase protein-encoding DNA or RNA in the cells and of activation of telomerase, which is also indicative of malignancy or an increased likelihood of progression to

-8-

malignancy. As described herein, the *hEST2* DNA encodes an RNA transcript which is expressed in primary tumors, cancer cell lines and telomerase - positive tissues and is readily detectable in post-crisis, telomerase-positive immortalized  
5 cells, but not detectable in telomerase-negative cell lines and differentiated telomerase-negative tissues. Detection of *hEST2* DNA and/or of the RNA transcript, thus, makes it possible to detect DNA which encodes a transcript required for telomerase activation that occurs during cellular  
10 immortalization and tumor progression and to detect the RNA transcript itself.

In an alternative embodiment, the invention is a method of diagnosing or aiding in the diagnosis of development of malignancy (cancer, tumor formation) in an  
15 individual, in which the occurrence (presence or absence) and/or quantity of a telomerase protein (e.g., *hEST2* protein) in cells is assessed. In the method, cells are obtained from an individual in need of diagnosis of malignancy and processed, if necessary, in such a manner  
20 that proteins in the cell are available for detection, such as by binding with antibodies or antibody fragments. The processed cells are combined or contacted with antibodies that recognize (bind) the telomerase protein (e.g., antibodies that bind Est2p or *hEST2* protein) under  
25 conditions appropriate for antibody binding to occur. Whether binding occurs is determined; optionally, the extent to which binding occurs can also be determined. If binding of anti-telomerase protein antibodies (e.g., anti-*hEST2* protein antibodies) to a component of processed cells  
30 occurs, it is indicative of the presence of *hEST2* protein in the cells and of malignancy or an increased likelihood of development or malignancy in the individual.

In one embodiment, the present invention is a method of reducing expression of *hEST2* RNA and *hEST2* protein in

- 9 -

cells of an individual (e.g., a human or other mammal).  
The method comprises administering to the individual a drug  
selected from the group consisting of drugs which inhibit  
(directly or indirectly) or bind *hEST2* RNA and prevent or  
5 reduce production of *hEST2* protein and drugs which inhibit  
*hEST2* protein function or activity. The drug is  
administered in a therapeutically effective amount (an  
amount sufficient to have the desired effect of reducing  
expression of *hEST2* RNA and *hEST2* protein) and under  
10 conditions appropriate for entry into cells, in which they  
have the desired effect.

In a further embodiment, the present invention is a  
method of treating cancer in an individual, in need of such  
treatment. In the method, a drug which inhibits or binds  
15 *hEST2* RNA (or DNA) and prevents or reduces production of  
*hEST2* protein or a drug which inhibits *hEST2* protein  
function or activity is administered to the individual  
under conditions appropriate for entry into cells and in a  
therapeutically effective amount (an amount sufficient to  
20 have the desired effect of inhibiting *hEST2* RNA and/or  
preventing or reducing production of *hEST2* protein or  
inhibiting *hEST2* protein function or activity), with the  
result that cancer in the individual is treated (reduced,  
reversed or prevented from advancing).

25 A further embodiment of the invention is a method of  
altering (increasing or reducing) lifespan of cells in  
culture or in an individual. In the method in which  
lifespan is increased, *hEST2* DNA is introduced into cells  
(e.g., in culture or in the individual), in which the *hEST2*  
30 RNA transcript and encoded protein are produced in  
sufficient quantity to increase the lifespan of cells in  
the cultured cells or the individual. In the method in  
which lifespan is decreased, *hEST2* protein function,

-10-

activity or production is reduced (partially or totally). This is done, for example, by introducing into cells in culture or administering to an individual in whom cell lifespan is to be decreased a drug which inhibits or binds  
5 *hEST2* RNA and prevents or reduces production of *hEST2* protein; a drug which inhibits *hEST2* protein function, activity or production; or a drug which inhibits or binds *hEST2* DNA and prevents or reduces production of *hEST2* RNA transcripts. The drug is administered to an individual  
10 under conditions appropriate for entry into cells in sufficient quantity to have the desired effect (in sufficient quantity to decrease lifespan of the cells by reducing *hEST2* protein function, activity or production).

In the above-described embodiments, the drug can be,  
15 for example, a small organic molecule, an enzyme which degrades *hEST2* protein, an enzyme inhibitor (e.g., a telomerase or *hEST2* enzyme inhibitor), an *hEST2* transcriptional regulator; an antisense molecule or a dominant negative form of *hEST2* protein.

20 Also described herein is a mutant yeast strain, referred to as DN, which carries an inactive chromosomal *rad52Δ* allele. The *RAD 52* coding sequence of a haploid strain of genotype *MATa leu 2-3, 112 lys2-201 trp1-1 ura3-52 his3-200* was replaced with a selectable marker (e.g.,  
25 the *HIS3* gene), to generate strain DN. Mutant yeast strain DNR, in which the absence of *RAD52* was complemented by introducing the plasmid *pYPCR*<sup>35</sup>, which encodes the *RAD52* gene under its own promoter and the selectable marker *URA3*, is also described. Also described are mutant yeast strain  
30 *DNRt1cΔ*, in which the *TLC1* gene in DNR cells was replaced with the *LEU2* selectable marker gene and Y0035, in which

-11-

the Est2 gene has been mutated with the addition of a TN3 LEU2 transposon in the DNR genetic background.

The mutant yeast strain described herein requires telomerase activity for viability. It is useful to carry  
5 out a rapid and automatable biological assay for telomerase inhibitors and, thus, for inhibitors of telomere biosynthesis. The assay is also the subject of the present invention. The mutant yeast strain is also useful to identify agents which are recombination inhibitors. In the  
10 mutant yeast strain, referred to as DN, the *RAD52* recombination gene has been replaced by a *HIS3* gene. As described herein, disruption of the *TLC1* telomerase RNA subunit gene or the *MIT1* putative telomerase protein subunit gene in the *rad52* mutants leads to a delayed cell  
15 death. Yeast lacking both recombination activity and telomerase activity die after about 20 to 60 cell generations. Telomerase activity can be disrupted by enzyme inhibitors, as well as by genetic inactivation. Thus, for a telomerase inhibitor screen, test compounds are  
20 added, at different concentrations, to the DNA mutant strain (the test strain) and to a wild-type (control) strain. Periodically, such as each day, cultures of the two strains are diluted (e.g., 1:1000). Compounds that kill the yeast instantly or that kill the DNA and wild-type  
25 yeast with identical delays, are eliminated. Compounds that kill the test yeast strain significantly before the control yeast strain are telomerase inhibitors. This biological assay has several advantages over available methods. It is fast, automatable, and can be performed  
30 easily in large quantities. Because it is a biological assay, it reflects the physiological function of the drugs. Furthermore, it allows testing of biochemical and uptake parameters in a single step. Finally, this assay also

-12-

excludes inhibitors of other metabolic enzymes, such as RNA and DNA polymerases and is specific for telomerase inhibitors.

#### BRIEF DESCRIPTION OF THE DRAWINGS

5        Figures 1A and 1B present the nucleic acid sequence of the *EST2* gene (SEQ ID NO.: 1) and the predicted amino acid sequence of the Est2 protein (SEQ ID NO. 2).

Figure 2 shows alignment of the predicted amino acid sequence of hEST2, also referred to as *hTERT*, (SEQ ID NO.: 10 3) with the yeast Est2p (SEQ ID NO.: 2) and Euplotes p123 (SEQ ID NO.: 4) homologues. Amino residues within shaded blocks are identical between at least two proteins. Identical amino acids within the reverse transcriptase (RT) motifs (Poch et al., *EMBO J.* 8:3867-3874 (1989); Xiong and 15 Eickbush, *EMBO J.* 9:3353-3362 (1990)) are in black boxes, an example of a telomerase-specific motif in an outlined gray box, and all other identical amino acids in gray boxes. RT motifs are extended in some cases to include other adjacent invariant or conserved amino acids. The 20 sequence of the expressed sequence tag AA281296 is underlined.

Figures 3A-3F show the alignment of RT motifs 1 to 6 of telomerase subunits hEST2 (SEQ ID NO.: 5-10), p123 (SEQ ID NO.: 11-16) and Est2p (SEQ ID NO.: 17-22) with S. 25 cerevisiae group II intron-encoded RTs a2-Sc (SEQ ID NO.: 23-28) and a1-Sc (SEQ ID NO.: 29-34). The consensus sequence of each RT motif is shown (h: hydrophobic residues, p: small polar residues, c: charged residues). Amino acids that are invariant among telomerases and the RT 30 consensus are boxed in black, while those that are invariant among telomerases and similar to the RT consensus are boxed in gray. Open boxes identify highly conserved

-13-

residues unique to either telomerases or to non-telomerase RTs. Asterisks denote amino acids essential for polymerase catalytic function.

Figure 4 is an ideogram of human chromosome 5p showing linkage of *hEST2* to sequence-tagged site (STS) markers WI-9907 and D5S417 determined by radiation hybrid (RH) mapping employing the Genebridge 4 RH panel (GB) (1 cR=270 kb). The higher resolution Stanford G3 RH panel (G3) placed *hEST2* close to marker D5S678 (1 cR=30 kb). The calculated distance between *hEST2* and these STS markers is in centiRays (cR) with LOD scores in parentheses.

Figures 5A-5B show the nucleotide sequence of *hEST2* cDNA in which the start codon ATG is underlined (SEQ ID NO.: 35).

Figure 6 is the amino acid sequence of *hEST2* (SEQ ID NO.: 3).

Figures 7A, 7B and 7C are the nucleotide sequence of partially or alternatively spliced *hEST2* message RNA (SEQ ID NO.: 36).

Figures 8A-8F show the expression of *hEST2* in normal human tissues and cancer cell lines. Duplicate RNA blots were probed with an *hEST2* probe (top panels) or with a  $\beta$ -actin probe as internal control (bottom panels). The 4.4 kb, 6 kb, and 9.5 kb transcripts are *hEST2*-specific.

Figures 9A-9D shows *hEST2* expression in primary human tumors. Rnase protection assays are shown for *hEST2* and  $\beta$ -actin controls. Sizes of the full-length and protected bands are indicated. Shown are the HL60 leukemia cell line (control), normal breast tissue, 2 primary breast tumors, the MCF7 and T47D breast cancer cell lines, and normal and primary tumor tissues from the testis, colon, and ovary. The doublet seen protected by the *hEST2* probe is invariant, and may be a result of probe secondary structure.

-14-

## DETAILED DESCRIPTION OF THE INVENTION

The linear chromosomes of eukaryotic cells offer the biological advantages of rapid recombination, assortment, and genetic diversification. However, linear DNA is inherently more unstable than circular forms. To address this difficulty, the eukaryotic chromosome has evolved to include a DNA-protein structure, the telomere, which caps chromosome ends and protects them from degradation and end-to-end fusion (Blackburn, *Ann. Rev. Biochem.* 53:163-194 (1984); Zakian, *Science* 270:1601-1607 (1995)).

The DNA component of telomeres consists of tandem repeats of guanine-rich sequences that are essential for telomere function (Blackburn, *Ann. Rev. Biochem.* 53:163-194 (1984), Blackburn, *Nature* 350:569-573 (1991); Zakian, *Science* 270:1601-1607 (1995)). These repeats are replicated by conventional DNA polymerases and by a specialized enzyme, telomerase (Greider, In *Telomeres*, eds. Blackburn and Greider, Cold Spring Harbor Laboratory Press, pp. 35-68 (1995)), first identified in the ciliate *Tetrahymena* (Greider and Blackburn, *Cell* 43:405-413 (1985)). The telomerase enzyme is essential for complete replication of telomeric DNA because the cellular DNA-dependent DNA polymerases are unable to replicate the ultimate ends of the telomeres due to their requirement for a 5' RNA primer and their unidirectional mode of synthesis. Removal of the most terminal RNA primer following priming of DNA synthesis leaves a gap that cannot be replicated by these polymerases (Olovnikov, *Dokl. Akad. Nauk SSSR* 201:1496-1499 (1971); Watson, *Nature New Biol.* 239:197-201 (1972)). Telomerase surmounts this problem by *de novo* addition of single-stranded telomeric DNA to the ends of chromosomes (Greider and Blackburn, *Cell* 43:405-413 (1985)),



-15-

Greider and Blackburn, *Nature* 337:331-337 (1989); Yu et al., *Nature* 344:126-132 (1990); Greider, In *Telomeres*, eds. Blackburn and Greider, Cold Spring Harbor Laboratory Press, pp. 35-68 (1995)).

5       The telomerase enzymes that have been characterized to date are RNA-dependent DNA polymerases which synthesize the telomeric DNA repeats using an RNA template that exists as a subunit of the telomerase holoenzyme (Greider, In *Telomeres*, eds. Blackburn and Greider, Cold Spring Harbor Laboratory Press, pp. 35-68 (1995)). The genes specifying the RNA subunits of telomerases have been cloned from a wide variety of species, including humans (Feng et al., *Science* 269:1236-1241 (1995); Greider, In *Telomeres*, eds. Blackburn and Greider, Cold Spring Harbor Laboratory Press, pp. 35-68 (1995)) and been shown in several instances to be essential for telomerase function *in vivo* (Yu et al., *Nature* 344:126-132 (1990); Yu and Blackburn, *Cell* 67:823-832 (1991); Singer and Gottschling, *Science* 266:404-409 (1994); Cohn and Blackburn, *Science* 269:396-400 (1995);

20   McEachern and Blackburn, *Nature* 376:403-409 (1995)). In addition, three proteins have been identified to date that are associated with telomerase activity. p80 and p95 were purified from the ciliate *Tetrahymena* (Collins et al., *Cell* 81:677-686 (1995)) and the gene encoding a mammalian

25   homologue of p80, TP1/TLP1, has also been cloned (Harrington et al., *Science* 275:973-977 (1997); Nakayama et al., *Cell* 88:875-884 (1997)). The specific mechanism by which these proteins participate in telomerase function has not been defined.

30       Perturbing the function of telomerase in the ciliate *Tetrahymena*, through the overexpression of an inactive form

-16-

of the telomerase RNA, or in yeast, through the mutation of genes encoding either the catalytic protein or template RNA subunit, leads to progressive telomere shortening as cells pass through successive cycles of replication (Yu et al.,  
5 *Nature* 344:126-132 (1990); Singer and Gottschling, *Science* 266:404-409 (1994); McEachern and Blackburn, *Nature* 376:403-409 (1995); Lendvay et al., *Genetics* 144:1399-1412 (1996); Counter et al., *Proc. Natl. Acad. Sci., USA* 94:9202-9207 (1997); Lingner et al., *Science* 276:561-567  
10 (1997)). This loss of telomeric DNA is ultimately lethal if it is not overcome. The lethality seems to be triggered when telomeres have been truncated below a critical threshold level. Hence, in the absence of compensating mechanisms, yeast cell lineages that lack telomerase  
15 activity have a lifespan dictated by the lengths of their telomeres.

In humans, telomerase activity is readily detectable in germline cells and in certain stem cell compartments. However, enzyme activity is not detectable in most somatic  
20 cell lineages (Harley et al., *Cold Spring Harb. Symp. Quant. Biol.* 59:307-315 (1994); Kim et al., *Science* 266:2011-2015 (1994); Broccoli et al., *Proc. Natl. Acad. Sci., USA* 92:9082-9086 (1995); Counter et al., *Blood* 85:2315-2320 (1995); Hiyama et al., *J. Immunol.* 155:3711-  
25 3715 (1995)). Consistent with this, telomeres of most types of human somatic cells shorten with increasing organismic age and with repeated passaging in culture, similar to the situation seen in protozoan and yeast cells that have been deprived experimentally of a functional  
30 telomerase enzyme (Harley et al., *Nature* 345:454-460 (1990); Hastie et al., *Nature* 346:866-868 (1990)).

-17-

Eventually the proliferation of cultured human cells will halt at a point termed senescence (Hayflick and Moorhead, *Exp. Cell. Res.* 25:585-621 (1961); Goldstein, *Science* 249:1129-1133 (1990)), apparently before the telomeres of  
5 these cells have become critically short.

Cultured normal human cells can circumvent senescence and thereby continue to proliferate when transformed by a variety of agents. In such cultures, telomere shortening continues until a subsequent point is reached that is  
10 termed crisis, where telomeres have become extremely short (Counter et al., *EMBO J.* 11:1921-1929 (1992) and *J. Virol.* 68:3410-3414 (1994); Shay et al., *Oncogene* 8:1407-1413 (1993); Klingelhutz et al., *Mol. Cell. Biol.* 14:961-969 (1994). Crisis, perhaps best described in SV-40  
15 transformed cells, is characterized by karyotypic instability, particularly the types of instability observed in chromosomes lacking functional telomeres, and by significant levels of cell death (Sack, *In Vitro* 17:1-19 (1981)). The crisis phenotype is reminiscent of that  
20 observed in yeast and Tetrahymena cells in which telomerase function has been experimentally perturbed.

The simplest interpretation of these data is that the lifespan of telomerase-negative human cells, like that of their yeast and ciliate counterparts, is ultimately limited  
25 by the length of telomeres. Rare human cells that have acquired the ability to grow indefinitely emerge from crisis populations with a frequency of  $10^{-6}$  to  $10^{-7}$  (Huschtscha and Holliday, *J. Cell Sci.* 63:77-99 (1983); Shay and Wright, *Exp. Cell Res.* 184:109-118 (1989)). This  
30 implies that a mutational event is required to confer the immortal phenotype on these cells. The immortal cells that escape crisis are characterized by readily detectable

-18-

levels of telomerase activity and by stable telomeres (Counter et al., *EMBO J.* 11:1921-1929 (1992) and *J. Virol.* 68:3410-3414 (1994); Shay et al., *Mol. Cell. Biol.* 15:425-432 (1995); Whitaker et al., *Oncogene* 11:971-976 (1995);  
5 Gollahon and Shay, *Oncogene* 12:715-725 (1996); Klingelhutz et al., *Nature* 380:79-82 (1996)). This suggests that activation of telomerase can overcome the limitations imposed by telomere length on the lifespan of cell lineages.

10       Activation of telomerase also appears to be a major step in the progression of human cancers. Unlike normal human cells, cancer cells can be established as permanent cell lines and, thus, are presumed to have undergone immortalization during the process of tumorigenesis.  
15 Moreover, telomerase activity is readily detected in the great majority of human tumor samples analyzed to date (Counter et al., *Proc. Natl. Acad. Sci., USA* 91:2900-2904 (1994); Kim et al., *Science*, 266: 2011-2015 (1994); Shay and Bacchetti, *Eur. J. Cancer* 33:787-791 (1997)).

20       Taken together, these various observations have been incorporated into a model which proposes that the limitation on prolonged cell replication imposed by telomere shortening serves as an important anti-neoplastic mechanism used by the body to block the expansion of  
25 pre-cancerous cell clones. According to such a model, tumor cells transcend the crisis barrier and emerge as immortalized cell populations by activating previously unexpressed telomerase, enabling them to restore and maintain the integrity of their telomeres (Counter et al.,  
30 *EMBO J.* 11:1921-1929 (1992) and *Proc. Natl. Acad. Sci., USA* 91:2900-2904 (1994); Harley et al., *Cold Spring Harb. Symp. Quant. Biol.* 59:307-315 (1994)).

-19-

A major question provoked by this model is the mechanism used to resurrect telomerase expression during tumor progression. Expression of the telomerase-associated protein TP1/TLP1 does not reflect the level of telomerase activity (Harrington et al., *Science* 275:973-977 (1997); Nakayama et al., *Cell* 88:875-884 (1997)). It is also clear that the levels of the human telomerase RNA component, *hTR*, cannot completely explain the regulation of telomerase activity. Although the levels of *hTR* and its mouse counterpart, *mTR*, increase with tumor progression (Feng et al., *Science* 269:1236-1241 (1995); Blasco et al., *Nat. Genet.* 12:200-204 (1996); Broccoli et al., *Mol. Cell. Biol.* 16:3765-3772 (1996); Soder et al., *Oncogene* 14:1013-1021 (1997)), the amounts of these transcripts do not always correlate with enzymatic activity. Indeed, *hTR* or *mTR* transcript levels can be significantly higher in telomerase-negative cells and tissues than in telomerase-positive cancer cells (Avilion et al., *Cancer Res.* 56:645-650 (1996); Bestilny et al., *Cancer Res.* 56:3796-3802 (1996); Blasco et al., *Nat. Genet.* 12:200-204 (1996)). Similarly, even though telomerase levels increase 100- to 2000-fold during the immortalization of human cells, the level of *hTR* message increases at most two-fold (Avilion et al., *Cancer Res.* 56:645-650 (1996)). Therefore, derepression of the *hTR* and TP1 subunits cannot easily be invoked to explain the appearance of telomerase activity in the great majority of human tumor samples. Thus far, the rate-limiting step in telomerase activation has remained elusive.

Described herein is work which has shown that mutations in a gene of eukaryotic origin (in an open

-20-

reading frame required for telomere maintenance) results in loss (absence) of telomerase enzymatic activity in cells and that the encoded protein is physically associated with the active telomerase enzyme, thus showing that the DNA  
5 encodes a component of a eukaryotic telomerase holoenzyme. Also described herein is isolated DNA of eukaryotic origin which is required for telomerase enzymatic activity and the isolated encoded RNA transcripts and proteins. Specifically, *EST2*, a yeast gene, and *hEST2*, which is of  
10 human origin and the encoded respective transcripts and proteins (yeast *EST2p* and human *hEST2*) are described, as are their respective sequences. DNA of the present invention includes cDNA, genomic DNA, recombinantly produced DNA and chemically synthesized DNA. It can be  
15 obtained from a source in which it occurs in nature or can be produced using known chemical or recombinant DNA methods. The following is a discussion of work carried out relating to yeast and human telomerase protein.

Gene Encoding Protein Subunit Of Yeast Telomerase  
20 Holoenzyme and the Encoded Protein

As described herein, it has been shown, using a genetic screen in budding yeast, that mutations in *EST2*, an open reading frame required for telomere maintenance, (Lendvay, T.S. et al., *Genetics* 144, 1399-1412 (1996))  
25 result in the absence of telomerase enzymatic activity in cells. As also described herein, epitope tagging and immunochemical studies showed that the *Est2* protein is physically associated with the active telomerase enzyme. Thus, *EST2* encodes a component of the yeast telomerase  
30 holoenzyme.

Interest in telomeres and telomerase has heightened in recent years with the discovery that telomerase is present

-21-

at low, almost undetectable, levels in most human somatic tissues and is readily detectable in germline cells and in the vast majority of tumor cell samples (Counter, C.M. et al., *Proc. Natl. Acad. Sci. USA* 91, 2900-4 (1994); Kim, N.W. et al., *Science* 266, 2011-5 (1994). Somatic cell lineages which lack telomerase lose telomere segments progressively as they pass through successive replication cycles, limiting their lifespan (Harley, C.B. et al., *Nature* 345, 458-60 (1990), Hastie, N.D. et al., *Nature* 346, 866-8 (1990)). Conversely, cell populations that resurrect telomerase expression ensure maintenance of telomere length, thereby removing a barrier to their further unlimited replication and resulting in the process termed immortalization (Counter, C.M. et al., *Embo. J.* 11, 1921-9 (1992)). These observations have led to a model which proposes that the process of telomere shortening limits the replicative potential of most human somatic cell lineages, and that cancer cells overcome this limitation by activating telomerase expression during the course of tumor progression (Counter, C.M. et al., *Embo. J.* 11, 1921-9 (1992); Harley, C.B. et al., *Cold Spring Harb. Symp. Quant. Biol.* 59, 307-15 (1994)).

Validation of this model has been held back by the difficulties associated with isolating and characterizing the telomerase enzyme and its encoding genes. Much of this work has focused on the telomerase enzymes of ciliates; the formation of thousands of mini-chromosomes present in their macro-nuclei seems to require correspondingly high amounts of telomerase, dwarfing the amounts available for study in metazoan cells. The telomerase enzyme in *Tetrahymena* has been reported to be composed of two proteinaceous and one RNA subunit, the latter responsible for templating the

-22-

- telomeric DNA segments (Greider, C.W. & Blackburn, E.H., *Nature* 377, 331-7 (1989); Collins, K. et al., *Cell* 81, 677-86 (1995); Yu, G.L. et al., *Nature* 344, 126-32 (1990)). The protein subunits of telomerase have not been described
- 5 in any other organism, although the genes specifying the RNA subunits of telomerase have been cloned in a wide range of other species (Blasco, M.A. et al., *Science* 269, 1967-70 (1995); Feng, J., et al., *Science* 269, 1236-41 (1995); Lingner, J. et al., *Genes Dev.* 8, 1984-98 (1994);
- 10 McEachern, M.J. & Blackburn, E.H., *Nature* 376, 403-9 (1995); Melek, M. et al., *Mol. Cell Biol.* 14, 7827-38 (1994); Shippen-Lentz, D. & Blackburn, E.H., *Science* 247, 546-52 (1990); Singer, M.S. & Gottschling, D.E., *Science* 266, 404-9 (1994)).
- 15 Genetic strategies designed to identify and isolate telomerase-encoding genes have centered on the yeast, *S. cerevisiae*. Mutations in several genes have been found to lead to progressive telomere shortening. Of these genes, *TLC1*, which specifies the RNA subunit of the telomerase
- 20 enzyme (Singer, M.S. & Gottschling, D.E., *Science* 266, 404-9 (1994)), is required for telomerase enzymatic activity (Cohn, M. & Blackburn, E.H., *Science* 269, 396-400 (1995); Lin, J.J. & Zakian, V.A., *Cell* 81, 1127-35 (1995); Lue, N.F. & Wang, J.C., *J. Biol. Chem.* 270, 21453-6 (1995);
- 25 Steiner, B.R. et al., *Proc. Natl. Acad. Sci. USA* 93, 3817-21 (1996)). On the other hand, *EST2* encodes a protein of unknown function (Lendvay, T.S. et al., *Genetics* 144, 1399-1412 (1996)); *EST1* (Lundblad, V. & Szostak, J.W., *Cell* 57, 633-43 (1989)) and *CDC13* (Garvik, B. et al., *Mol. Cell*
- 30 *Biol.* 15, 6128-38 (1995); Nugent, C.I., Hughes, T.R., Lue,



-23-

N.F. & Lundblad, V., *Science* 274, 249-52 (1996)) are not essential for telomerase activity (Cohn, M. & Blackburn, E.H., *Science* 269, 396-400 (1995); Nugent, C.I. et al., *Science* 274, 249-52 (1996)); and *KEM1* appears to encode a  
5 nuclease that acts on telomeres (Liu, Z. & Gilbert, W., *Cell* 77, 1083-92 (1994); Liu, Z. et al., *Proc. Natl. Acad. Sci. USA* 92, 6002-6 (1995)). A search for genes encoding telomerase protein subunits was initiated because the protein subunits of the yeast telomerase have not yet been  
10 identified, and because there are no obvious homologs of the *Tetrahymena* telomerase protein subunits encoded in the yeast genome (Collins, K. et al., *Cell* 81, 677-86 (1995); Cherry, J.M. et al., *Saccharomyces Genome Database* (<http://genome-www.stanford.edu/Saccharomyces>)).

15 A genetic screen designed to identify the genes specifying telomerase subunits was undertaken, based on the observation that telomere loss in yeast cells lacking the telomerase RNA component can be compensated by the actions of a second telomere-maintenance system that utilizes the  
20 DNA recombination machinery. An essential component of this machinery is the product of the *RAD52* gene. Accordingly, the simultaneous inactivation of *RAD52* and the telomerase RNA component results in cell lethality in both *S. cerevisiae* and *K. lactis* (Lendvay, T.S. et al., *Genetics*  
25 144, 1399-1412 (1996); McEachern, M.J. & Blackburn, E.H., *Genes Dev.* 10, 1822-34 (1996)). This lethality is only observed after some delay and occurs in concert with the extensive telomere shortening observed following serial passaging of the mutant cells. Applicants reasoned that  
30 mutation of other genes contributing to telomerase function, including the genes encoding the protein

-24-

subunit(s) of the enzyme, might be recognized because they similarly render the yeast dependent on wild-type *RAD52* for prolonged viability. If so, telomerase mutant yeast could be identified by their requirement for *RAD52* function.

5       A yeast strain carrying an inactive chromosomal *rad52Δ* allele that was complemented by a plasmid-borne wild-type *RAD52* gene linked to a *URA3* gene was generated, in order to determine if mutants in telomerase function could be revealed by a screen for mutants that require *RAD52*  
10 function. When *RAD52* function was inactivated through eviction of the *RAD52*-encoding plasmid (by counter-selecting against the *URA3* marker with 5-fluoro-orotic acid (FOA), cell viability was unaffected for the succeeding 80 cell generations (Boeke, J.D. et al., *Mol. Gen. Genet.* 197,  
15 345-6 (1984). However, when the *TLC1* gene, which encodes the RNA subunit of telomerase, was additionally inactivated in this strain, a decrease in cell viability was observed after 60 generations. Moreover, when the rescuing *RAD52* plasmid was evicted, there was a marginal decrease in  
20 viability after 40 generations and complete inviability after 60 generations. This showed that telomerase-deficient cells perish upon the loss of *RAD52* function and that this phenotype is therefore useful to screen for mutants carrying lesions in a variety of genes affecting  
25 telomerase function.

      The *rad52Δ* mutant cells carrying the complementing *RAD52* plasmid were mutagenized by homologous recombination with a library of yeast genomic fragments, each of which carried one or more copies of an inserted mini-  
30 *Tn3::lacZ::LEU2* transposon (Burns, N., et al., *Genes Dev.* 8, 1087-105 (1994)). To ensure a high density of mutations,

-25-

1x10<sup>6</sup> transposon-mutagenized yeast clones were generated. These mutagenized yeast cells were cultured for 40 generations to allow the telomeres of any cells lacking telomerase function to shorten substantially. A portion of  
5 the cell colonies was replica-plated and the *RAD52* plasmid was evicted from one of each replica pair to identify cell clones that required *RAD52* for continued viability.

Of the 10<sup>5</sup> transposon-mutagenized clones plated after 40 generations, approximately 2,500 reproducibly lost  
10 viability following removal of the *RAD52* plasmid. In this case, as earlier, these cells were placed in the presence of FOA to select against the presence of the plasmid-borne *URA3* gene. The loss of viability following FOA might have derived through a second, unrelated mechanism in which the  
15 *URA3* gene became stabilized through chromosomal integration, rendering cells FOA-sensitive independent of any effects on telomeres. For this reason, the FOA-sensitive cell clones were mated with wild-type yeast, yielding diploids; all cell clones for which the diploid  
20 derivatives remained FOA-sensitive were discarded.

The 245 remaining *RAD52*-dependent cell clones were tested for changes in telomere structure by Southern hybridization analysis of their telomeres. Of these 245 haploids, 16 cell clones carried short telomeres identical  
25 to those seen in yeast lacking a functional *TLC1* gene. Tetrad analysis of the diploid derivatives of these clones was performed to determine whether the *rad52* synthetic lethal phenotype was caused by the transposon insertion or by an adventitious unlinked mutation elsewhere in the yeast  
30 genome. In each case, the lethality co-segregated with the transposon insertion, demonstrating that this phenotype is

-26-

derived directly from the genetic disruption effected by a single inserted transposon.

To determine which genes were disrupted by transposon insertion in the 16 yeast strains containing short  
5 telomeres, the transposons together with flanking genomic DNAs from each strain were cloned and the DNA sequences flanking the transposon insertion site were analyzed. This sequencing revealed that 6 clones represented 6 distinct insertion sites within the *TLC1* gene which, as mentioned  
10 above, specifies the RNA subunit of the telomerase. This result provided strong and direct validation of the utility of the strategy described to screen for genes that directly affect telomerase function.

Eight of the cell clones having shortened telomeres  
15 and lethality in the absence of *RAD52* carried transposons that had inserted into seven distinct positions within an uncharacterized open reading frame on chromosome XII at nucleotides 766540 to 769194 (Cherry, J.M. et al., *Saccharomyces* Genome Database (<http://genome-www.stanford.edu/Saccharomyces>)).  
20

The predicted amino acid sequence encoded by the gene identified on chromosome XII is presented in Figures 1A and 1B (SEQ ID NO.: 2); the predicted protein includes 884 amino acids with an estimated mass of 102 kilodaltons. The  
25 predicted gene product does not share significant sequence similarity with any sequences in the databases available through the National Center for Biotechnology Information BLAST server. However, subsequent to completion of the work described herein, a report was published on the  
30 identification of a novel gene, *EST2*, involved in telomere length regulation. The open reading frame identified on chromosome XII is identical to *EST2*; therefore, this gene is referred to herein as *EST2*.

-27-

Genetic analysis of the *est2* transposon-insertion mutants shows that they have the same delayed cell death phenotype, dependence on *RAD52*, and telomere shortening as was displayed by cells carrying mutations in their *TLC1* gene. Diploids heterozygous for disrupted alleles of *RAD52* and *EST2* were sporulated and the resulting tetrads were analyzed for growth in culture and for telomere length. In 33 tetrads analyzed, the vast majority of the *est2::TN3 rad52Δ* double mutant spore products became inviable by approximately 20 generations in culture, and the remainder died by approximately 40 generations. The viability of the *est2::Tn3* mutants with wild-type *RAD52* began to decline by approximately 60 generations, and this phenotype was more pronounced after approximately 80 generations. The *EST2* wild-type spore products remained completely viable at all generations, regardless of their *RAD52* genotype. Consistent with the role of *EST2* in telomere maintenance, the decrease in telomere length, by which this gene was identified, co-segregated with the transposon insertional mutations in *EST2*. This telomere shortening was comparable to that observed in *tlc1::TN3* mutants.

Until now, *TLC1* has been the only yeast gene known to be a subunit of telomerase and to be required for telomere activity *in vitro*. The *est2::TN3* gene mutants are phenotypically indistinguishable from *tlc1::TN3* mutants. Analyses were carried out to determine whether *EST2* is also required for telomerase function.

A telomerase assay was used to show that extracts from wild-type yeast catalyze the elongation of a single-stranded telomeric primer by four to sixteen nucleotides in an RNase-sensitive fashion, indicating telomerase activity.

-28-

However, extracts derived from yeast bearing transposons inserted into *EST2* lacked detectable telomerase activity, like *tlc1Δ* mutants, which are known to be telomerase-negative (Cohn, M. & Blackburn, E.H., *Science* 269, 396-400  
5 (1995)). Comparison with dilutions of wild-type extract made it possible to estimate that extracts from both *tlc1* and *est2* mutants contain at most 5% of wild-type telomerase activity. While the actual levels of telomerase in the mutant yeast are likely to be much lower, such lower levels  
10 could not be quantified with this assay. As expected from the phenotypic analysis of tetrads, the absence of detectable telomerase activity co-segregated with transposon insertion, independent of the *RAD52* allele.

The requirement of *EST2* function for telomerase  
15 activity suggests that Est2 is either a protein subunit of telomerase or an upstream regulator required for telomerase activity. It was possible to distinguish between these possibilities by replacing the endogenous *EST2* gene with a variant encoding a protein with three influenza  
20 hemagglutinin (HA) epitope stages at its carboxy terminus (*EST2-HA*). This modified *EST2* allele and, thus, its product, is fully functional. The *EST2-HA* yeast show no overt phenotype, their telomeres are wild-type length, and extracts from these yeast exhibit normal levels of  
25 telomerase activity. Extracts from *EST2-HA* and control yeast were incubated with excess anti-HA antibody, after which the immunoprecipitates and resultant supernatants were separated and assayed for telomerase activity. The anti-HA antibody immunoprecipitated telomerase activity  
30 from extracts of yeast expressing the tagged Est2-HA protein, but not from extracts expressing normal Est2 protein. The immunoprecipitation was specific for the

-29-

anti-HA antibody since a control anti-Myc epitope antibody failed to immunoprecipitate activity from either extract. Essentially all of the telomerase activity remained in the supernatant of the anti-HA immunoprecipitation from wild-type extracts, whereas incubation with the anti-HA antibody almost completely immunodepleted the telomerase activity from the Est2-HA extracts. These data indicated that Est2 is not only physically associated with telomerase, but is a constituent of most, if not all, of the active telomerase complex.

Intriguingly, *EST2* shares no homology with either of the two genes reported to encode the *Tetrahymena* telomerase protein subunits (Collins, K. et al., *Cell* 81, 677-86 (1995)). A search of the yeast genome (Cherry, J.M. et al., *Saccharomyces Genome Database* (<http://genome-www.stanford.edu/Saccharomyces>) revealed no open reading frame sharing obvious sequence similarity with the *Tetrahymena* genes. Thus, the genes encoding the yeast and ciliate telomerase enzymes may be highly diverged, the Est2 function may be absent in *Tetrahymena*, or the analogous Est2 protein of *Tetrahymena* may not yet have been identified.

Work described herein supports the essential role of *EST2* gene for telomerase activity *in vitro* and *in vivo*. Immunochemical experiments show that Est2-containing complexes contain telomerase activity, and that removal of these complexes from the cell concomitantly removes the telomerase activity. These characteristics argue that Est2 is an essential and integral component of the yeast telomerase holoenzyme. The identification of this protein subunit of yeast telomerase should aid in the characterization of the other elements of this complex,

-30-

both in yeast and in other species, as well as in the understanding of the biochemical mechanism by which this enzyme acts.

Work described herein provides the basis for a method  
5 of assessing the ability of a molecule or compound to act as a telomerase inhibitor or as a recombination inhibitor. It provides a method by which a drug (molecule or compound) which inhibits telomerase or inhibits recombination can be identified. Eukaryotic linear chromosomes are capped with  
10 telomeres, a DNA-protein complex that protects the integrity of the chromosome end. Telomeric DNA sequences are composed of multiple repeat elements. When the telomeric DNA is lost from chromosome ends, cells die because of the resulting chromosomal instability.

15 Telomere length is maintained by two mechanisms. In almost all eukaryotes growing under normal conditions, telomeric DNA repeats are synthesized by the ribonucleoprotein enzyme telomerase. This RNA-dependent DNA polymerase (or "reverse transcriptase") copies the DNA  
20 repeats from an RNA template that is a component of the enzyme.

When the telomerase pathway is inactive, however, telomere length can be maintained by an alternative mechanism in yeast and possibly also in humans: the  
25 homologous recombination pathway used for double strand break repair.

In the yeast *Saccharomyces cerevisiae*, mutants of the telomerase subunits in conjunction with mutants of the recombination pathway have been shown to become inviable  
30 after growth for 40 to 80 generations. This synthetic lethal phenotype was used as a screen for telomerase mutants in a background of recombination mutants and, as a result, a novel telomerase gene was identified.



-31-

Drug screens which work in the same fashion as demonstrated herein for the genetic screen are useful to identify telomerase inhibitors in a recombination mutant background, and recombination inhibitors in a telomerase mutant background.

Telomere maintenance inhibitors, which can be a telomerase inhibitor alone, a recombination inhibitor alone, or a combination, can be identified by the method described herein, using appropriate mutant eukaryotic cells, such as the mutant yeast strain described herein. Such telomerase inhibitors can be further assessed, using known methods, to confirm their effectiveness as anti-fungal drugs or as anti-cancer drugs. Normal human somatic cells lack a pathway to maintain telomere lengths, and their telomeres shorten continuously. Immortal human cells, whether in tissue culture or the vast majority of malignant tumors in the human body, have developed a mechanism to maintain telomeres. Telomere maintenance occurs as a result of activation of the telomerase enzyme, activation of a recombination pathway, or possibly both. In yeast, the inactivation of both pathways has been shown to kill cells; in immortal tumor cells, pharmacological inhibitors must be tested before the effects of inactivation of either or both pathways can be assessed. A brief description of screens for telomerase inhibitors and recombination inhibitors follows.

#### Screen for Telomerase Inhibitors

As described below, a mutant strain of the yeast *Saccharomyces cerevisiae* that requires telomerase activity for viability has been generated. This strain can be used in a rapid and automatable biological assay for telomerase inhibitors.

-32-

In the mutant yeast strain, named DN, the *RAD52* recombination gene has been replaced by a *HIS3* gene. Disruption of the *TLC1* telomerase RNA subunit gene or the *MIT1/EST2* telomerase protein subunit gene in the *rad52* mutants has been shown to lead to a delayed cell death. Yeast lacking both recombination activity and telomerase activity die after about 20 to 60 cell generations.

Telomerase activity can be disrupted by enzyme inhibitors as well as by genetic inactivation, and, therefore, the mutant strains described can be used to identify telomerase inhibitors. For a telomerase inhibitor screen, test compounds are added at different concentrations to the DN mutant strain and to a wild-type control strain. Each day, the cultures are be diluted (e.g., 1:1000). Compounds that kill the yeast instantly, or that kill the DN and wild-type yeast with identical delays, are eliminated. Compounds that kill the DN yeast significantly before the wild-type yeast are telomerase inhibitors.

The method described is a biological assay and has several advantages over available methods. It is fast, automatable, and can be performed easily in large quantities. As a biological assay, it reflects the physiological function of the drugs. Furthermore, it allows testing of biochemical and uptake parameters in a single step. Finally, this assay will also exclude inhibitors of other metabolic enzymes, such as RNA and DNA polymerases and is highly specific for telomerase inhibitors.

High-throughput screening can be accomplished by growing the yeast in microtiter dishes. Aliquoting of media, dilution of test drug compounds, and dilution of yeast cultures can be performed robotically. Growth can be

-33-

measured by the optical density of the yeast culture at 595-600 nm; growth measurements and preliminary analysis of the data can also be automated.

The telomerase inhibitor screen has several  
5 applications, such as to identify inhibitors of fungal telomerase enzymes, mammalian telomerase enzymes and reverse transcriptase.

1) Inhibitors of fungal telomerase enzymes. This screen is a rapid method for identification of compounds that  
10 inhibit telomerase in the yeast *Saccharomyces cerevisiae*. This yeast is closely related to the fungal pathogen *Candida albicans* and, therefore telomerase inhibitors effective in *Saccharomyces* will likely function in *Candida*, as well as in other pathogenic yeast. There are only a few  
15 effective anti-fungal drugs currently available; many are inadequate because of their serious side effects. Thus, a novel approach to anti-fungal drugs could be clinically valuable for the treatment of systemic fungal infections.

2) Inhibitors of mammalian telomerase enzymes. Telomerase  
20 function is likely to be conserved between diverse species. Therefore, it is highly possible that inhibitors of yeast telomerase could also inhibit mammalian telomerase, and can be tested for their ability to act as anti-cancer agents. These compounds can also serve as lead compounds for the  
25 identification of such inhibitors and for modification to produce more effective inhibitors (e.g., inhibitors which are longer-lived, exhibit great inhibitory effects, resist degradation by cellular enzymes).

3) Inhibitors of reverse transcriptase. The drugs that  
30 have been identified as telomerase inhibitors to date have been shown to be inhibitors of HIV reverse transcriptase. Telomerase inhibitors identified by the present screen can

-34-

be assessed, using known methods, for their ability to act as retroviral reverse transcriptase.

#### Screen for recombination inhibitors

In this case, inhibitors of the telomere recombination pathway are identified by their ability to kill telomerase mutant yeast but not wild-type yeast. The screen is similar to the telomerase inhibitor screen described above.

For a recombination inhibitor screen, test compounds are added at different concentrations to a telomerase mutant strain (e.g., *tlc1* or *mit1/est2*), referred to as a test strain, and to a wild-type control strain. Each day, the cultures are diluted 1:1000. Compounds that kill the yeast instantly, or that kill the telomerase mutant and wild-type yeast with identical delays, are eliminated.

Compounds that kill the telomerase mutant yeast significantly before the wild-type yeast are recombination inhibitors. The ability of the compounds to inhibit recombination can be confirmed using known methods. The recombination inhibitors have several uses. For example, combination therapy of a recombination inhibitor with a telomerase for treating fungal infections and/or malignant neoplasms. In addition, recombination inhibitors can be used to potentiate the toxic effects of radiation treatment and of chemotherapeutic agents that induce DNA double-strand breaks.

The following methods and materials were used in the work described herein.

#### METHODS

Yeast strains: Strain DN was generated by replacing the *RAD52* coding sequence of the haploid strain L3853 (gift of G. Fink; genotype *MATa leu2-3,112 lys2-201 trp1-1 ura3-*

-35-

52 *his3-200*) with the *HIS3* gene using a PCR-based homologous gene disruption method (Baudin, A. et al., *Nucleic Acids Res.* 21, 3329-30 (1993)). The absence of *RAD52* in this strain was complemented by introducing the  
5 plasmid pYPCR<sup>35</sup>, which encodes the *RAD52* gene under its own promoter and the selectable marker *URA3*, thereby generating strain DNR. The *TLC1* gene was replaced with the *LEU2* selectable marker by the above PCR procedure in L3853 and DNR cells, yielding the strains *tlc1Δ* and *DNRtlc1Δ*,  
10 respectively.

Yeast mutagenesis: Approximately 230 μg of the appropriately prepared yeast::mini-Tn3::lacZ::*LEU2* genomic library (Burns, N., et al., *Genes Dev.* 8, 1087-105 (1994)) were introduced by homologous recombination into  
15 approximately 7x10<sup>6</sup> cells of strain DNR. Approximately 1x10<sup>6</sup> Ura<sup>+</sup>Leu<sup>+</sup> clones representing successful transposon insertions were pooled and replated to a total number of ~1x10<sup>6</sup> clones to "age" the yeast for the *rad52Δ* synthetic lethal screen. 1x10<sup>5</sup> of such yeast were then reseeded for  
20 the *rad52Δ* synthetic lethal screen, as described in the text. DNA flanking insertion sites were rescued as described by Burns, sequenced and compared to the *Saccharomyces* Genome Database (Stanford University School of Medicine, Department of Genetics) (Burns, N., et al.,  
25 *Genes Dev.* 8, 1087-105 (1994)).

Southern hybridization: Genomic DNA was isolated and digested with the restriction enzyme *XhoI*, which liberates the telomeric DNA (Guthrie, C., & Fink, G.R., *Guide to Yeast Genetics and Molecular Biology* (Academic Press, new  
30 York, 1991); Chan, C.S. & Tye, B.K., *Cell* 33, 563-73

-36-

(1983)). 2  $\mu$ g of this DNA was resolved on 0.8% agarose gel for 570 Vh, and Southern hybridized as described by Counter et al. using a  $^{32}$ P kinase labelled yeast telomeric oligonucleotide CACCACACCCACACACCACA (SEQ ID NO.: 37)  
5 (Counter, C.M., et al., *Embo. J.* 11, 1921-9 (1992)).

Tetrad analysis: Diploid derivatives of the transposon-mutagenized yeast lacking the pYCPR+ plasmid were sporulated and tetrads were dissected according to standard methods (Guthrie, C., & Fink, G.R., *Guide to yeast*  
10 *genetics and molecular biology* (Academic Press, New York, 1991)). Presence of the *rad52::HIS3* allele and the *miniTn3::lacZ::LEU2* insertions were assayed on -His and -Leu synthetic complete plates, respectively.

Yeast extracts: 6 litres of yeast cultures were  
15 harvested at an optical density of 0.4-0.6 at 600 nm, resuspended in TMG buffer, disrupted with a bead beater and centrifuged for 90' at 100,000g at 4°C as described (Cohn, M. & Blackburn, E.H., *Science* 269, 396-400 (1995)). The supernatant was decanted, flash frozen and stored at -70°C.  
20 Extracts typically had a protein concentration of 16-20 mg/ml.

Telomerase assay: Telomerase activity was assayed essentially as outlined by Cohn and Blackburn with the following two modifications: first, 4  $\mu$ g of crude S100  
25 yeast extract was incubated with the oligonucleotide TGTCTGGGTGTCTGGG (SEQ ID NO.: 38) and second, the reaction products were purified and resolved on 15% acrylamide, 7M urea sequencing gels (Counter, C.M. et al., *Proc. Natl. Acad. Sci. USA* 91, 2900-4 (1994)).

30 Epitope tagging: An HA-URA3-HA cassette was introduced into the 3' end of the *EST2* gene by homologous recombination, and then counter-selected on FOA to excise

-37-

the *URA3* gene and generate three copies of the influenza hemagglutinin tag in frame with *EST2* (Schneider, B.L. et al., *Yeast* 11, 1265-74 (1995)). The construction of this epitope-tagged version deleted the 10 carboxy-terminal amino acids of *Est2*; these codons contain a TA repeat sequence that is present in multiple copies in the yeast genome.

Immunoprecipitation and immunodepletion: 540  $\mu$ g of yeast extract was diluted in 170  $\mu$ l of TMG buffer (Cohn, M. & Blackburn, E.H., *Science* 269, 396-400 (1995)) supplemented with 150 mM NaCl (buffer TMGN) and incubated at 4°C with 20  $\mu$ l of packed protein A-agarose beads pre-loaded with anti-HA antibody (monoclonal antibody 12CA5). After 2 hours, the mixture was briefly centrifuged to separate the agarose beads from the supernatant extract. The beads were then washed four times with buffer TMGN and resuspended in 50  $\mu$ l TMGN. As a control, a duplicate extract was diluted and left untreated on ice for 2 hours. 4  $\mu$ g of supernatant or untreated extracts and 10  $\mu$ l of beads were assayed for telomerase activity.

#### Human Telomerase Catalytic Subunit Gene and Encoded Protein

The findings presented herein support modulation of *hEST2* (*hTERT*) RNA expression levels as an important mechanism used in a variety of developmental contexts to determine the amount of telomerase activity present in specific cell lineages. Indeed, the levels of the catalytic subunit of telomerase may represent a rate-limiting determinant of enzyme activity in many types of cells. Moreover, up-regulation of *hEST2* message may be an important mechanism through which telomerase becomes activated during both cellular immortalization and the progression of malignant tumors.

-38-

hEST2, (also referred to as hTERT) the human telomerase subunit described here, shares extensive sequence similarities with the catalytic subunits of the yeast and ciliate telomerase enzymes. The amino acid  
5 sequence conservation between these three enzymes is extensive and is scattered throughout their reading frames. In addition, no related genes were detected in the human genome by Southern blotting analysis. Together, these data support the conclusion that the gene described here encodes  
10 the catalytic subunit of the human telomerase holoenzyme. Further proof that this is the case can be obtained using known methods, such as experiments analogous to those performed with the yeast *EST2* gene, described herein and by Lingner and co-workers. (Lingner et al., *Science* 276:561-  
15 567 (1997)). For example, experiments that will provide additional proof that the gene described herein encodes the human telomerase subunit include those that demonstrate that mice lacking the mouse *mEST2* homologue also lack telomerase activity; that the hEST2 protein is physically  
20 associated with a ribonucleoprotein complex that exhibits telomerase activity; and that alteration of critical residues in the domain of hEST2 that is homologous to RTs inactivates its catalytic function.

As described herein, three telomerases described to  
25 date (hEst2, yeast Est2 and Euplotes p123) include motifs that indicate they are distant homologues of a variety of RTs. They also share several unique, telomerase-specific sequence motifs. Yeast, ciliates and mammals represent highly diverged branches of the phylogenetic tree, which  
30 suggests that the catalytic subunit of the telomerase was developed early in eukaryotic evolution. It is possible that it was present in the cell that became ancestral to all contemporary eukaryotes. Even earlier, it appears that



-39-

this enzyme shares ancestry with the precursors of the RTs specified by a variety of transposons and viruses.

The repression of *hEST2* mRNA in telomerase-negative cells and tissues and its upregulation that is associated with a number of human tumor cell lines and primary human tumors suggest one mechanism by which telomerase activity might be modulated. During development, the expression of the *hEST2* mRNA may be repressed in many post-embryonic cell lineages, depriving cells in these lineages of the telomerase catalytic subunit. This in turn may underlie the observed progressive telomeric shortening associated with aging in the cells in many of these lineages.

In contrast, the reappearance of telomerase enzyme activity when transformed cells escape from crisis or when tumors progress toward malignancy may, in many cases, be explained mechanistically by the de-repression of *hEST2* mRNA expression. As shown here, in cell cultures, this de-repression occurs in both transformed embryonic kidney cells and lymphocytes when they emerge from crisis and begin to exhibit telomerase activity. Not addressed here is whether the enhanced *hEST2* RNA expression is achieved at the transcriptional or post-transcriptional level.

Described here is a correlation between *hEST2* mRNA levels and assayable telomerase activity. These two manifestations of *hEST2* gene expression are present in a constant, predictable ratio. This provides the basis for assessing cells for *hEST2* mRNA as an indicator of telomerase activity or assessing cells for telomerase activity as an indicator of *hEST2* mRNA occurrence or level. It is possible that other mechanisms besides the presently observed modulation of *hEST2* mRNA levels may intervene to modulate telomerase activity.

-40-

As described in Example 8, over expression of *hEST2* in previously (normally) telomerase-negative cells is sufficient to impart telomerase activity to these cells, providing clear proof that *hEST2* is the human telomerase catalytic subunit. The data demonstrate that the ectopic expression of *hTERT* in otherwise telomerase-negative human cells is necessary and sufficient for induction of telomerase activity. In addition, the physical association of *hTERT* with telomerase activity confirms that *hTERT* is a telomerase subunit. Up-regulation of the *hTERT* gene is the sole barrier to activation of telomerase in the tested cells.

The identification of *hEST2* as the gene specifying the telomerase catalytic subunit provides an entree into understanding one of the essential steps in human tumor pathogenesis -- that leading to cell immortalization. Enzymatic assays have demonstrated that telomerase is often activated at a relatively late step in tumor progression (Harley et al., *Nature* 345:458-460 (1994)). This step may occur when evolving, pre-malignant cell clones have surmounted the senescence barrier and subsequently, having exhausted their telomeric ends, encounter the successive barrier of crisis. At this stage, the activation of telomerase may enable cells to breach these barriers to further clonal expansion, thereby conferring great selective advantage on the rare cell that has acquired the ability to resurrect the long-repressed telomerase activity. Indeed, activation of telomerase may represent an essential step in tumor progression.

Such dependence on telomerase activity means that this enzyme represents an attractive target of drugs designed to interfere with malignant cell proliferation. The finding that *hEST2* message is up-regulated in human tumors and in

-41-

immortalized cells lends further credence to this idea. Furthermore, the identification of hEST2 as the candidate telomerase catalytic subunit provides a biochemical reagent for identifying such drugs.

5 Described herein is a human gene, *hEST2*, which is a telomerase catalytic subunit gene. The gene transcript (mRNA) is expressed in primary human tumors, cancer cell lines and telomerase-positive tissues, but is undetectable, using the assays described herein, in cell lines known to  
10 be telomerase negative and in differentiated telomerase-negative tissues. The *hEST2* gene shares significant sequence similarity with a yeast telomerase catalytic subunit gene (*S. cerevisiae* EST2) and a ciliate telomerase catalytic subunit gene (*Euplotes* p123), as shown in Figure  
15 2. The three telomerase enzymes are members of the reverse transcriptase family of enzymes. There are seven conserved sequence motifs which define the polymerase domains of reverse transcriptases and six of these domains are present in *hEST2*, including the invariant aspartic acid residues  
20 which are required for telomerase activity. Also the subject of the present invention is DNA which encodes *hEST2* protein of SEQ ID NO.: 3; DNA which is the complement of DNA of SEQ ID NO.: 35; DNA which hybridizes to the complement of DNA of SEQ ID NO.: 35; DNA which hybridizes  
25 to the complement of DNA of SEQ ID NO.: 35 and encodes *hEST2* protein of SEQ ID NO.: 3, DNA which is a gene which localizes to human chromosome 5, subband 5p15.33. A RNA transcript or RNA message encoded by *hEST2* DNA of SEQ ID NO.: 35 or by DNA which encodes *hEST2* protein of SEQ ID  
30 NO.: 3 is also the subject of this invention.

As shown in Figure 2, conceptual translation of the 4030 bp *hEST2* DNA shows an open reading frame of 1132 amino acids which is predicted to be an approximately 127 kDa

-42-

protein. As also shown in Figure 2, there is a clear relatedness among hEST2, yeast EST2 and Euplotes p123 in regions of the gene outside the RT motifs, particularly sequence identities in a region just before motif 1 of the hEST2 protein. This region of similarity appears unique to telomerases and, thus, can be used to produce reagents, such as antibodies reactive with the unique region and nucleic acid probes and primers useful to identify hEST2 DNA, using, for example, hybridization or amplification methods. hEST2 is a single-copy gene which is approximately 40 kb in size and was shown to localize (to be present on) chromosome 5, particularly subband 5p15.33. (See Figure 4) Expression levels in normal human tissues and human cancer cell lines were assessed, as described in Example 3. This assessment revealed two major RNA species (4.4 kb and 9.5 kb) and one minor RNA species (approximately 6 kb). hEST message was detectable in several normal tissues, including thymus, testis and intestine; the latter two tissues are known to be telomerase-positive and the telomerase status of thymus has not been reported. Using the assays described herein, hEST2 transcript was not detected in most other normal human tissues, including heart, brain, placenta, liver, skeletal muscle and prostate, all of which have been reported to be telomerase-negative.

As described herein, the hEST2 transcript or message (mRNA) is expressed in primary human tumors and cancer cell lines, but is not detectable in telomerase-negative cell lines and differentiated (normal, nontumorigenic) telomerase-negative tissues. Further, hEST2 message is not detectable in pre-crisis, telomerase-negative transformed cells, but is readily detectable in post-crisis, telomerase-positive immortalized cells. The work described

-43-

herein provides a method of differentiating between telomerase-positive cells and telomerase-negative cells, as well as for monitoring a change in telomerase content of cells. As a result, as described below, the present work  
5 provides a method of identifying cells which are telomerase positive and, thus, of identifying cells which are tumor or cancer cells or are en route to becoming tumor cells (malignant).

A method of identifying telomerase-positive cells is  
10 the subject of the present invention. Also the subject of the present invention is a method of identifying cells which are transformed, malignant, cancerous, tumor or post-crisis cells or are en route to or likely to become transformed, malignant, cancerous, tumor or post-crisis  
15 cells. For convenience, the term malignant or malignancy is used herein to refer to transformed, cancer, cancerous, tumor, tumorigenic and post-crisis cells. For example, the phrase "a method of identifying malignant cells in an individual" encompasses a method of identifying cancer  
20 cells, transformed cells, tumor cells and/or post-crisis cells in an individual. The present invention, thus, provides a method of diagnosing, aiding in the diagnosis of or predicting an increased likelihood of the occurrence of cancer, tumor formation and/or the development of  
25 malignancy in an individual, particularly a mammal and, specifically, a human. In the method, telomerase content of cells is assessed by detecting or measuring *hEST2* DNA, *hEST2* transcript (*hEST2* message) or *hEST2*-encoded protein (*hEST2* protein). The quantity of *hEST2* DNA, *hEST2*  
30 transcript or *hEST2* protein can be determined; alternatively, the occurrence (presence or absence) of *hEST2* DNA, *hEST2* transcript or *hEST2* protein can be detected. The presence of *hEST2* DNA, *hEST2* message and/or

-44-

hEST2 protein in cells is indicative of the presence of malignant cells or malignancy in the individual.

In one embodiment, *hEST2* DNA, *hEST2* RNA (or both) is detected or measured in tissue or cells obtained from an individual. This can be carried out using known methods, such as hybridization (e.g., *in situ* hybridization) or amplification methods. All or a portion of *hEST2* DNA or RNA can be used in such a method as a probe (e.g., to detect *hEST2* DNA or RNA) or a primer (e.g., to amplify *hEST2* DNA).

As described herein, up-regulation of *hEST2* RNA is associated with activation of telomerase during cell immortalization and may be an important mechanism through which telomerase becomes activated during cellular immortalization, the progression of malignant tumors or both. Thus, detection and/or measurement of *hEST2* RNA provides a means of assessing cells (e.g., mammalian, such as human cells) for malignancy or the likelihood of progression to malignancy. For example, detection of *hEST2* RNA, even without quantification of *hEST2* RNA levels, in cells indicates that telomerase is activated and that the cells are malignant or have an increased likelihood of progression to malignancy (relative to cells in which *hEST2* RNA is not detected). Quantification of *hEST2* RNA can be carried out as well. Higher concentrations of *hEST2* RNA cells are expected to indicate a more advanced stage in malignancy or a greater likelihood of progression to malignancy than is the case with lower *hEST2* RNA levels in cells. *hEST2* RNA levels determined by assessing an individual's cells can be compared with pre-established *hEST2* RNA levels. For example, *hEST2* RNA levels determined

-45-

for an individual's cells can be compared with *hEST2* RNA levels in cells known to be malignant and/or at established stages of malignancy (e.g., a pre-established reference or standard).

5           Alternatively, the levels of *hEST2* RNA determined for an individual can be compared with *hEST2* RNA levels determined for the same individual in prior assessments, in which case the individual serves as his or her own standard or reference. In this approach, it is possible to monitor  
10 (assess over time) the status of the individual. This is useful, for example, to assess or aid in assessing whether the malignancy or cancer is progressing, remaining unchanged or regressing in that individual. This is particularly useful in assessing the status of an  
15 individual's malignancy or cancer before, during and after therapy (e.g., chemotherapy, radiation, surgery) and to assess the therapeutic value or suitability of a drug(s) or other agent(s) for a specific individual in need of treatment. As mentioned, assessment of *hEST2* RNA (or DNA)  
20 can be carried out using known methods, such as *in situ* hybridization, Rnase protection assay, reverse transcriptase (RT) PCR, Southern blot analysis or Northern blot analysis.

          In one embodiment, the presence of malignancy or an  
25 increased likelihood of malignancy is assessed in a method in which a nucleic acid (e.g., DNA) anneals to nucleic acid (e.g., RNA). For example, the method can be Northern blot analysis, Rnase protection assay or RT PCR. One embodiment is as follows: Cells to be assessed for *hEST2* DNA or RNA  
30 are obtained from the individual and processed to render DNA and/or RNA in the cells available for annealing with or hybridization of complementary nucleic acid sequences DNA or RNA (e.g., with complementary poly- or oligonucleotides

-46-

(DNA or RNA)). All or a portion of *hEST2* DNA or *hEST2* RNA can be used as the hybridization probe or sequence. As described herein, *hEST2* protein shares sequence similarities and identities with yeast p123 and Euplotes Est2p, particularly in a region just before motif 1 (Figure 2, boxed region). As described, this region appears to be unique to telomerases. Thus, it is useful in assays which identify and/or quantify telomerase in cells and provide an assessment of malignancy or the likelihood of progression to malignancy of cells (e.g., from an individual in whom a diagnosis of malignancy or assessment of the likelihood of the occurrence of malignancy is needed). DNA or RNA comprising DNA or RNA encoding region(s) of similarity and/or identify with yeast p123 and/or Euplotes Est2p such as region(s) unique to telomerase, can be used in the hybridization assay or other assay carried out to detect or measure *hEST2* DNA or RNA in an individual's cells.

Alternatively, DNA or RNA comprising DNA or RNA which encodes a region or regions unique to *hEST2* protein can be used as the probe. In an alternative approach, the *hEST2* DNA or RNA is used as a primer in an amplification method. The cells processed to render DNA and/or RNA available for annealing (hybridization) with complementary poly- or oligonucleotides (processed cells) are combined with all or a portion of *hEST2* DNA or RNA under conditions appropriate for hybridization of complementary nucleic acids (DNA, RNA) to occur. Whether annealing (hybridization) occurs is then determined. If hybridization occurs (forming complexes of cellular DNA or RNA and the *hEST2* poly- or oligonucleotides), it is an indication that the cells contain *hEST2* DNA or *hEST2* RNA. Poly- or oligonucleotides used in the method can be, for example, DNA which encodes the telomerase motif described herein (e.g., DNA which



-47-

encodes amino acid residues 556 to 565 of SEQ ID NO.: 3 or amino acid residues 560 to 565 of SEQ ID NO.: 3) or amino acid residues 1 to 50 of hEST2 protein or a portion thereof. If hEST2 RNA is upregulated (expressed), it is  
5 indicative of activation of telomerase and of malignancy or an increased likelihood of progression to malignancy. The amount of hEST2 RNA can also be determined (by determining, for example, the extent to which hybridization with hEST2 DNA or RNA occurs) and used to assess the extent of  
10 telomerase activity and the stage of malignancy. This detection and/or measurement of hEST2 DNA or RNA can be carried out at various intervals over time to assess the status of the malignancy (e.g., progression, reversal).

In another embodiment of assessing telomerase content  
15 and, thus, malignancy or cancer of cells, hEST2 protein is analyzed (detected and/or quantified). This can be done using known methods, such as enzymatic assays or immunoassays, which indicate hEST2 protein is present in cells assessed and, optionally, quantify the protein. For  
20 example, hEST2 telomerase activity can be assessed by determining whether extension of a telomeric primer occurs when a sample (e.g., cells, cell fractions or cell component(s)) from an individual is combined with the telomeric primer under conditions appropriate for hEST2  
25 protein telomerase activity and telomeric extension. Alternatively, antibodies which recognize (bind) hEST2 protein can be used to determine if the protein is present in cells or other sample obtained from an individual. In the method, cells are obtained from an individual and  
30 processed or treated to render proteins in the cells available for binding with antibodies, thus producing processed cells. The processed cells are combined or contacted with antibodies which bind hEST2 protein and

-48-

whether binding occurs between antibodies which bind hEST2 protein (hEST2 protein-binding antibodies) and protein in the cells is determined. If hEST2 protein-binding antibody/protein binding occurs (to form complexes), it is indicative of the presence of hEST2 protein in the cells and, thus, of malignancy or an increased likelihood of progression to malignancy in the individual (in cells of the individual). The antibodies can be monoclonal antibodies or polyclonal antibodies (e.g., polyclonal sera) and can be specific for (bind or recognize only) hEST2 protein or can be nonspecific for hEST2 protein (bind or recognize hEST2 protein and other protein(s)). Humanized antibodies can also be used. Antibodies which recognize a motif or epitope unique to hEST2 and/or to members of the class to which hEST2 belongs are particularly useful. For example, antibodies which specifically bind the telomerase motif, represented in Figure 2, which is common to hEST2 protein, yeast p123 protein and/or Euplotes EST2 can be produced, using known methods, and used to assess cells for hEST2 protein. Alternatively, antibodies which specifically bind a motif which is unique to hEST2 protein (e.g., all or a portion of amino acid residues 1 to 50 of hEST2 protein) can be used. As discussed above with reference to hEST2 DNA and hEST2 RNA, the assessment can be detection (determination of presence or absence) or measurement (quantification). The presence of hEST2 protein is indicative of malignancy or at least of an increased likelihood of progression to malignancy.

Antibodies which bind or recognize (specifically or non-specifically) hEST2 protein are also a subject of the present invention. Nucleic acid probes and primers (poly- or oligonucleotides) comprising hEST2 DNA or RNA are also the subject of this invention. The poly- or oligonucleotides will vary in length and need to be of

-49-

sufficient length to bind to and remain bound to *hEST2* DNA or *hEST2* RNA under the conditions used. They will generally be at least four to six bases in length and can comprise the entire *hEST2* DNA or *hEST2* RNA (alone or with  
5 additional non *hEST2* DNA or RNA). Preferably the probes or primers will hybridize to at least a characteristic portion of *hEST2* DNA or *hEST2* RNA (a portion which is present in members of the class to which *hEST2* belongs), thus making it possible to identify *hEST2* DNA or *hEST2* RNA  
10 substantially to the exclusion of other proteins. For example, a probe or primer can comprise DNA or RNA which encodes a characteristic motif or region of *hEST2* protein, such as the telomerase motif (e.g., amino acids 556 to 565 or 560 to 565) or the amino terminal amino acid sequence of  
15 *hEST2* protein not present in Est2p or p123. (See Figure 2). For example, a probe or primer encoding amino acid residues 1 to 50 (inclusive) of *hEST2* protein (SEQ ID NO.: 3) or a portion thereof can be used.

Methods of altering *hEST2* DNA transcription and  
20 expression, methods of altering *hEST2* protein function and methods of identifying agents which alter (enhance or reduce) transcription, expression or function are also the subject of this invention. Also the subject of the present invention are a method of increasing or shortening the  
25 lifespan of cells in culture, *ex vivo* or *in vivo*; agents or drugs (DNA, RNA, drugs, small organic molecules, enzymes, for example) useful for lengthening or shortening cell lifespan and methods of identifying agents which enhance lifespan. For example, one embodiment of a method of  
30 inhibiting *hEST2* protein comprises introducing into cells an agent which inhibits *hEST2* protein, directly or indirectly; as a result, function of the enzyme is

-50-

abolished or the enzyme is inactivated. hEST2 protein is inhibited directly, for example, by introducing into cells an agent which binds to or otherwise "ties up" hEST2 protein such that it is less active or inactive (cellular telomerase activity is reduced or eliminated).  
5 Alternatively, a drug or agent which inactivates hEST2 or inhibits hEST2 catalytic function (and concomitantly reduces or eliminates telomerase activity) by degrading it or preventing it from being produced can be introduced into  
10 cells in which hEST2 protein is to be inhibited. Such drugs or agents can be, for example, a small organic molecule or a dominant negative protein. Agents which prevent production of hEST2 message or its further processing (e.g., to produce DNA) or a dominant negative  
15 form of the telomerase protein can also be introduced into cells to inhibit hEST2 protein. Methods and agents which inhibit hEST2 catalytic function are useful for treatment of cancer and, thus, are useful as anti-cancer therapies.

If hEST2 DNA, hEST2 RNA or hEST2 protein is inhibited  
20 (partially or totally) in cells, lifespan of the cells will be shortened (shorter than it would be if the DNA, RNA or protein were not inhibited). For example, if a cell is malignant or more likely to progress to malignancy because of hEST2 function, an agent or drug which inhibits hEST2  
25 transcription or expression or hEST2 protein function in the cell will also shorten the lifespan of the cell because telomerase activity (and, thus, chromosomal extension or maintenance) will be inhibited. The agent is useful as a therapeutic for treating or preventing malignancy in  
30 individuals in whom malignant cells or cells with increased likelihood to progress to malignancy are present. These agents include, for example, enzyme (telomerase or hEST2 protein) inhibitors, enzymes which degrade hEST2 protein,

-51-

hEST2 transcriptional regulators, antisense molecules (e.g., DNA, RNA, PNA) and dominant negative mutant forms of hEST2. In a method of the present invention in which lifespan of cells is altered (increased or decreased) in an individual, a drug or agent is introduced into the individual in such a manner that it enters cells of the individual in sufficient quantity to have the desired effect (increase or decrease in cell lifespan). For example, a drug which prevents production of hEST2 transcript of hEST2 protein function (directly or indirectly) is introduced into an individual, using known methods, in sufficient quantities to enter cells, such as a tumor, precancerous or cancerous cells, whose lifespan is to be decreased (e.g., made shorter than would be the case in the absence of the drug). Alternatively, a drug which enhances cell lifespan can be introduced into an individual in such a manner that it enters cells in sufficient quantity to enhance hEST2 protein expression or prolong its activity (e.g., by blocking its degradation).

20 The present invention is also a method of enhancing or increasing the lifespan of cells in culture, in which telomerase activity or function is enhanced, such as by introducing hEST2 DNA or hEST2 protein into the cells or by activating an endogenous hEST2 gene. For example, the lifespan of normal human or other mammalian cells can be extended (immortalized human or other mammalian cells can be produced) and the resulting cells used for therapeutic purposes (e.g., grafting of tissue (such as skin) or of organs), screening or assay methods or production of proteins or other cellular products. For example, the lifespan of epithelial cells, keratinocytes or endothelial cells can be extended by introducing hEST2 DNA into the cells, in which the hEST2 DNA is expressed or introducing

-52-

hEST2 protein into the cells. The resulting cells with longer lifespan can be transplanted into or grafted onto an individual (e.g., as skin grafts, as systems for delivery of therapeutic proteins, such as hormones and enzymes), to whom they provide therapeutic benefit. In the method of extending the lifespan of cultured cells, cells whose lifespan is to be extended are cultured under conditions appropriate for their viability and hEST2 DNA or hEST2 protein is introduced into them. The resulting cells are maintained under conditions appropriate for expression of hEST2 DNA and/or activity of hEST2 protein, with the result that the lifespan of the cells is enhanced. At an appropriate time (e.g., after sufficient cells with enhanced lifespan are available) enhanced lifespan cells (such as keratinocytes) are transplanted into an individual (e.g., as a skin graft). cells.

The present invention is illustrated by the following examples, which are not intended to be limiting in any way.

#### EXAMPLE 1: Cloning of hEST2 (hTERT)

The expressed sequence tag database (dbEST) was searched for sequences related to the yeast protein Est2p (Lendvay et al., *Genetics* 144:1399-1412 (1996)) and the Euplotes protein p123 (Lingner et al., *Science* 276:561-567 (1997)), using the program TBLASTN and the server <http://www.ncbi.nlm.nih.gov/BLAST/>. This resulted in the identification of a homologous expressed sequence tag, Genbank accession # AA281296, which derived from Soares NbHTGBC cDNA clone 712562. DNA from clone plasmid 712562 was PCR-amplified with primers HT-1 (5' -AAGTTCCTGCACTGGCTGATGAG- 3') (SEQ ID NO.: 39) and HT-5 (5' -TCGTAGTTGAGCACGCTGAACAG- 3') (SEQ ID NO.: 40). The

-53-

resulting 377 bp fragment was used to probe 1 ZAP phage cDNA libraries derived from the human Jurkat T-cell lymphoma (Stratagene, La Jolla, CA) and human Nalm-6 pre-B cell leukemia cell lines (Weissbach et al., *J. Biol. Chem.* 5 269:20517-20521 (1994)). A total of seven independent clones were isolated from these two libraries. Three of these cDNA clones together with plasmid 712562 were sequenced completely in both directions. The remaining clones were sequenced in specific regions. Reprobing of 10 the Jurkat library with the 5'-most 500 bp region identified from the *hEST2* cDNA clones yielded one new clone containing an insert that overlapped with already determined sequences.

Rapid amplification of cDNA ends (RACE) was performed 15 by PCR amplifying testis Marathon-ready cDNA (Clontech, Palo Alto, CA) with flanking primer AP-1 (Clontech) and *hEST2* primer R0096 (5' -CAAGAAACCCACGGTCACTCGGTCCACGCG-3') (SEQ ID NO.: 41), and then re-amplifying with flanking primer AP-2 and *hEST2* primer R0098 (5' 20 -CAGCTCCTTCAGGCAGGACACCTGCGGG-3') (SEQ ID NO.: 42). The product was subcloned and sequenced in both directions. RACE was also performed on aliquoted pools of plasmid DNA from a human testis cDNA library (Reduced Complexity cDNA Analysis, Qingyun Liu and Fang Chen, unpublished results) 25 with primers HT-21 (5' -CAGGTGACACCACACAGAAA-3') (SEQ ID NO.: 43) or HT-22 (5' -TTCCAAGCAGCTCCAGAAA-3') (SEQ ID NO.: 44) and a vector primer, re-amplified with the vector primer and *hEST2* primers R0098 or R0097 (5' -CCTTCGGGGTCCACTAGCGTGTGGCGG) (SEQ ID NO.: 45), then 30 purified and sequenced.

DNA sequencing reactions were performed with the AmpliTaq FS Prism ready reaction cycle sequencing kit (Perkin-Elmer/ABI) and electrophoresed on a 373 A Stretch

-54-

ABI DNA sequencer. The resulting sequences were assembled into a contig using the program Sequencher 3.0 (Gene Codes Corporation, Ann Arbor, MI). The 1132 amino acid ORF identified in this contig was aligned with p123 and Est2p using the Pattern-Induced Multi-sequence Alignment program version 1.4 (R. F. Smith, Baylor College of Medicine & T. F. Smith, Boston University) using the server <http://dot.imgen.bcm.tmc.edu:9331/multi-align/multi-align.html>, with minor modifications to the final alignment.

10 BLASTP searches were performed against the non-redundant protein databases using the server <http://www.ncbi.nlm.nih.gov/BLAST/>.

#### EXAMPLE 2: Identification of the *hEST2* Gene

The sequences derived from the catalytic subunit of the *S. cerevisiae* telomerase, Est2p, and from the *Euplotes* p123 protein (Lendvay et al., *Genetics* 144:1399-1412 (1996); Counter et al., *Proc. Natl. Acad. Sci., USA* 94:9202-9207 (1997); Lingner et al., *Science* 276:561-567 (1997)), were used to identify a human homologue (Genbank accession # AA281296) in the National Center for Biotechnology Information expressed sequence tag database. As shown in Figure 2 (underlined sequence), the conceptual translation of this DNA sequence shows clear relatedness to both the yeast and ciliate telomerase sequences as evidenced by identical or similar amino acid residues scattered throughout the entire length of the expressed sequence tag.

25

The identified expressed sequence tag provided only a fragment of the putative human telomerase open reading frame. Therefore, a human Jurkat T-cell lymphoma and a human Nalm-6 pre-B cell leukemia cDNA library were screened

30



with a probe derived from the expressed sequence tag. Seven cDNA clones were retrieved. The resulting sequence was extended further in the 5' direction by rapid amplification of cDNA ends (RACE) on human testis cDNA and on an independently generated human testis cDNA library.

Assembly of the cDNA clones and RACE products, together with the clone containing the expressed sequence tag, resulted in a contiguous sequence spanning 4030 bp. Conceptual translation of this 4 kbp sequence reveals an open reading frame of 1132 amino acids which is predicted to encode a 127 kDa protein (Figure 2). Although we have not identified an in-frame upstream stop codon in the sequence, the first methionine identified by RACE (Figure 2) is a candidate for the translation start site, as the sequence of this putative translation initiation site, CCCGCCAUGC (SEQ ID NO.: 46), is similar to the consensus GCC(A/G)CCAUGG (SEQ ID NO.: 47) characteristic of translation initiation sites (Kozak, Cell 44:238-292 (1986)).

The predicted 127 kDa protein shares extensive sequence similarity with the entire sequences of the Euplotes and yeast telomerase subunits (Figure 2) and extends beyond the amino- and carboxyl-termini of these proteins. A BLAST search reveals that the probabilities of these similarities occurring by chance are  $1.3 \times 10^{-18}$  and  $3 \times 10^{-13}$ , respectively. By way of comparison, the probability of similarity between the yeast and Euplotes telomerases in a protein BLAST search is  $6.9 \times 10^{-6}$ . The human gene was initially named *hEST2* (human EST2 homologue) to reflect its clear relationship with the yeast gene, the first of these genes to be described. The current name for the gene is *hTERT*, which stands for human TELOmerase Reverse Transcriptase. *EST2* was named because of the

-56-

phenotype of ever shortening telomeres caused by its mutant alleles (Lendvay et al., *Genetics* 144:1399-1412 (1996)) and was later demonstrated to encode the yeast telomerase catalytic subunit (Counter et al., *Proc. Natl. Acad. Sci.*,  
5 USA 94:9202-9207 (1997); Lingner et al., *Science* 276:561-567 (1997)).

Like the yeast and ciliate telomerase proteins, hEST2 (*hTERT*) is a member of the reverse transcriptase (RT) family of enzymes (Figures 2 and 3A-3F). Seven conserved sequence  
10 motifs, which define the polymerase domains of these enzymes, are shared among the otherwise highly divergent RT family (Poch et al., *EMBO J.* 8:3867-3874 (1989); Xiong and Eickbush, *EMBO J.* 9:3353-3362 (1990)). p123 and Est2p  
15 share six of these motifs with, most prominently, the a2-Sc enzyme, an RT that is encoded within the second intron of the yeast *COX1* gene (Kennell et al., *Cell* 73:133-146 (1993)). These six motifs, including the invariant aspartic acid residues known to be required for telomerase enzymatic function (Counter et al., *Proc. Natl. Acad. Sci.*,  
20 USA 94:9202-9207 (1997); Lingner et al., *Science* 276:561-567 (1997)), are found at the appropriate positions of the predicted sequence of hEST2 (Figures 2 and 3A-3F). Thus, the proposed human telomerase catalytic subunit, like its yeast and ciliate counterparts, belongs to the RT  
25 superfamily of enzymes.

Although hEST2 shares some sequence similarity with RTs, it is not a conventional RT. Rather, it is far more closely related to the telomerase catalytic subunits of yeast and ciliates than to other RTs. Whereas the BLAST  
30 probability of sequence similarity between hEST2 and the telomerase subunits of the single-cell eukaryotes arising by chance is  $10^{-13}$  to  $10^{-18}$ , the chance probability of

-57-

sequence similarity with the next most closely related RT, a2-Sc, is 0.12. Beyond the motifs that define the polymerase domains of these various enzymes, hEST2 shows no sequence similarity with RTs. In contrast, in its domains  
5 that lie N-terminal to the polymerase domain, BLAST searches identify clear relatedness between hEST2 and both p123 and Est2p, the chance occurrence of these similarities being  $1.6 \times 10^{-9}$  and  $1.8 \times 10^{-4}$  respectively. Many of the sequence identities in the N-termini of hEST2, p123 and  
10 Est2p reside in a region just before motif 1 (Figure 2, boxed region). This sequence is not found in RTs, nor is it apparent in other proteins, suggesting the presence of motifs that may be unique to telomerases. Identification of the catalytic subunits of yet other telomerases will be  
15 required to validate this possibility.

Even within the hEST2 domain that share sequence similarity with RTs, it is clear that hEST2 is more closely related to the already described telomerases than it is to non-telomerase RTs. For example, the sequence similarity  
20 of the region encompassing the RT motifs between hEST2 and the catalytic subunits of yeast and Euplotes has a probability of chance occurrence of  $5.7 \times 10^{-6}$  and  $1.9 \times 10^{-5}$  respectively compared to 0.0056 for a2-Sc, the next most closely related non-telomerase RT. Within the RT  
25 motifs are several amino acids that are invariant among the telomerases but divergent between telomerases and non-telomerase RTs, or alternatively nearly invariant among non-telomerase RTs but divergent between these RTs and telomerases (Figures 3A-3F).

30 In summary, hEST2, Est2p, and p123 form a clearly defined subgroup within the RT family. For these reasons, hEST2 is a human homologue and very likely an orthologue of the microbial enzymes described to date.

-58-

Sequencing of a number of cDNA clones has revealed two distinct forms of *hEST2* transcripts. See Figures 5A-5B and Figures 7A-7C. Four independent cDNA clones, isolated from three independently generated libraries deriving from  
5 distinct cell types, lack an identical 182 bp segment within the open reading frame. The absence of this segment leads to a shift in reading frame that introduces a premature termination codon. Both forms of the *hEST2* transcript were detected by RT-PCR in a variety of human  
10 cell types. Information on the intron-exon boundaries of *hEST2* is not available. The simplest interpretation of these data is that the sequence difference between the two groups of cDNAs reflects the existence of two alternatively spliced mRNAs of the *hEST2* gene. The physiological  
15 consequences of the expression of the potential non-functional *hEST2* transcript are obscure at present.

### EXAMPLE 3: Chromosomal Localization of *hEST2*

The *hEST2* cDNA was used as a probe in Southern blot analyses of human genomic DNA. These reveal *hEST2* to be a  
20 single-copy gene with an estimated size of 40 kb. All the genomic sequences reactive with the cDNA probe appear to be components of this ~40 kb locus, suggesting that there are no other closely related genes in the human genome.

*hEST2* was localized to a specific chromosomal region  
25 by analyzing two independent panels of hamster-human radiation hybrid (RH) cells with two markers spanning different regions of *hEST2*. Initial mapping using the Genebridge 4 RH panel placed both *hEST2* markers between sequence-tagged sites (STS) WI-9907 and D5S417 (Figure 4).  
30 Independent confirmation of this localization was then obtained by mapping carried out with a second panel, the

-59-

Stanford G3 RH panel. This second mapping placed both *hEST2* markers next to STS marker AFMA139YA9 (GDB locus D5S678) which itself is localized between the above-mentioned markers WI-9907 and D5S417. These markers  
5 are present at the telomeric end of chromosome 5p (Hudson et al., *Science* 270:1945-1954 (1995)).

Mapping of the *hEST2* locus was further refined by localizing it to subband 5p15.33, since the STS markers D5S678 and D5S417 have been assigned to this band on the  
10 Genethon YAC contig map (Chumakov et al., *Nature* 377:175-297 (1995)). This localization is consistent with fluorescence in situ hybridization (FISH) analysis of YAC 767E1. This YAC maps further away from the telomere than D5S678 and has been assigned to chromosome 5p15.33 by FISH  
15 (Chumakov et al., *Nature* 377:175-297 (1995)).

Chromosome 5p is one of the most common targets for amplification in non-small-cell lung cancers; it is amplified in ~70% of tumors (Balsara et al., *Cancer Res.* 56:645-650 (1997); Petersen et al., *Cancer Res.* 57:2331-  
20 2335 (1997)). The effects of this amplification on *hEST2* expression levels are unknown.

#### EXAMPLE 4: Expression of *hEST2* Transcripts

Telomerase becomes activated during tumor progression. and, as discussed above, this activation has been  
25 associated with the immortalization of tumor cells. A variety of mechanisms might be invoked to explain such activation, among which is the induction of the expression of one or more telomerase subunits. The telomerase holoenzyme is presumed to exist as a multi-subunit  
30 ribonucleoprotein complex and, therefore, the levels of any one of the subunits, including those of the catalytic

-60-

subunit described here, might be rate-limiting in determining enzyme activity. Alternatively, the components of the telomerase holoenzyme might be expressed constitutively and subject to various types of

5 post-translational modification that would govern their activity. As mentioned earlier, transcript levels of the telomerase RNA subunit and of the *TP1/TLP1* gene do not necessarily correlate with telomerase activity (Feng et al., *Science* 269:1236-1241 (1995); Avilion et al., *Cancer*

10 *Res.* 56:645-650 (1996); Blasco et al., *Nat. Genet.* 12:200-204 (1996); Harrington et al., *Science* 275:973-977 (1997); Nakayama et al., *Cell* 88:875-884 (1997)), implying that at least one other mechanism is responsible for regulating human telomerase.

15 The expression levels of *hEST2* mRNA in various cell types were analyzed, using both RNA Northern hybridizations and RNase protection assays, in order to address one of the remaining possible regulatory mechanisms,. RNA blots prepared from a panel of normal human tissues and human

20 cancer cell lines were probed with cDNA fragments deriving from two independent, non-overlapping regions of the *hEST2* gene. This probing revealed two major RNA species migrating near the 4.4 kb and the 9.5 kb markers, as well as a minor species of ~6 kb (Figures 8A-8F). Each of these

25 RNA species was recognized by both probes, confirming that each represents an *hEST2* mRNA.

*hEST2* message was detectable in several normal tissues including thymus, testis, and intestine. Of these, testis (Kim et al., *Science* 266:2011-2015 (1994); Wright et al.,

30 *Dev. Genet.* 18:173-179 (1996)) and intestine (Hiyama et al., *Int. J. Oncol.* 9:453-458 (1996)) are known to be

-61-

telomerase-positive, while the telomerase status of the thymus has not been reported. In marked contrast, the *hEST2* transcript was undetectable in our assays in most other normal human tissues, including heart, brain, placenta, liver, skeletal muscle, and prostate (Figures 8A, 8B, 8D and 8E), all of which have been reported to be telomerase-negative (Kim et al., *Science* 266:2011-2015 (1994); Wright et al., *Dev. Genet.* 18:173-179 (1996); Shay and Bacchetti, *Eur. J. Cancer* 33:787-791 (1997)). The absence of detectable *hEST2* message in ovary may seem paradoxical, as ovary is a germline tissue, and germline tissues have been reported to harbor significant levels of telomerase. However, oocyte division is completed by birth, and while fetal ovary is telomerase-positive (Kim et al., *Science* 266:2011-2015 (1994); Wright et al., *Dev. Genet.* 18:173-179 (1996)), both adult ovarian epithelium and mature oocytes are telomerase-negative (Counter et al., *Proc. Natl. Acad. Sci., USA* 91:2900-2904 (1994); Wright et al., *Dev. Genet.* 18:173-179 (1996)). It is also possible that the levels of *hEST2* are below the threshold of detection. The 7 kb band seen in muscle tissue appears to be an artifact, as it was not observed with an independent *hEST2* probe.

In contrast to its absence in the majority of normal tissues, *hEST2* mRNA was strongly expressed in a variety of cancer cell lines, most strikingly in the leukemic cell lines HL-60 and K-562 (Figures 8C and 8F). It is unclear why only very low levels of *hEST2* message were observed in HeLa cells on this particular Northern blot (Figures 8C and 8F); reanalysis of independently prepared HeLa cell RNA both by Northern blot and by RNase protection demonstrated

-62-

that *hEST2* mRNA is present in HeLa cells at high levels comparable to those seen in K-562 cells. The two major and one minor *hEST2* transcript appear to be expressed in the same relative proportions in all cell types that yielded  
5 detectable *hEST2* mRNA.

EXAMPLE 5: *hEST2* Expression in Primary Human Cancers

The expression of *hEST2* mRNA in cancer cell lines suggested that *hEST2* transcript levels might be elevated in primary tumors as well. To determine if this is the case,  
10 RNA was extracted from a variety of tumor samples as well as from normal control tissues and analyzed this for the presence of *hEST2* mRNA using an RNase protection assay. In total, 11 of 11 tumor samples examined showed detectable levels of *hEST2* message (Figures 9A-9D). *hEST2* RNA was  
15 undetectable in normal breast and ovarian tissue but was expressed at significant levels in 2 of 2 breast tumors as well as 2 of 2 breast tumor-derived cell lines, and in 4 of 4 ovarian tumors.

As expected from Northern hybridization analysis of  
20 *hEST2* RNA (Figures 8A-8F) and the known pattern of telomerase catalytic activity in human tissues (Kim et al., *Science* 266:2011-2015 (1994); Hiyama et al., *Int. J. Oncol.* 9:453-458 (1996)), the *hEST2* transcript was detected at high levels by the RNase protection assay in testis and at  
25 moderate levels in colon. Similarly, using the RNase protection assay, four colon tumor samples and a testicular tumor sample were found to express detectable levels of *hEST2* RNA (Figures 9A-9D); two of the colon tumors showed significantly elevated levels as well. These data suggest  
30 that the *hEST2* message is expressed in a very high



-63-

percentage of tumors and is specifically induced in tumors that arise from telomerase-negative tissues.

EXAMPLE 6: Up-regulation of *hEST2* mRNA Is Associated with  
Telomerase Activation and Cellular Immortalization

5       The strong expression of the *hEST2* message observed in several tumors and cancer cell lines suggested that the levels of this transcript are correlated with the amount of telomerase enzyme activity in these various cell types. To investigate this possibility, telomerase activity and *hEST2*  
10 RNA level were analyzed in a panel of non-immortalized and immortalized cell lines. Using the TRAP telomerase assay (Kim et al., *Science* 266:2011-2015 (1994)), two mortal fibroblast strains, WI-38 and IMR-90, were found to lack detectable telomerase activity. In contrast, telomerase  
15 activity was readily detectable in three immortal cell lines, HeLa, 293, and K-562. The telomerase-negative cells also lacked detectable *hEST2* message as gauged by an RNase protection assay, while the immortal telomerase-positive cell lines expressed significant levels of *hEST2* RNA.  
20 These results indicate that *hEST2* message levels correlate closely with telomerase activity.

      Thus, as described herein, *hEST2* RNA expression and telomerase activity are present in immortal transformed cells, but absent in mortal normal cells. These findings  
25 support the role of induction of *hEST2* expression in the activation of telomerase that occurs during cellular immortalization. That this is the case was shown by analyzing *hEST2* transcript levels by RNase protection, and comparing the levels of hTR and telomerase activity in  
30 pre-crisis cells prior to the up-regulation of telomerase, and in post-crisis telomerase-positive immortal cells from

- 64 -

two different transformed human cell populations:  
Epstein-Barr virus-transformed B lymphocytes (B4 cells) and  
SV40-T antigen transformed embryonic kidney cells (HA1  
cells). Result observed were consistent with previously  
5 reported assessments (Counter et al., *EMBO J.* 11:1921-1929  
(1992), *Proc. Natl. Acad. Sci., USA* 91:2900-2904 (1994);  
Avilion et al., *Cancer Res.* 56:645-650 (1996)). Little or  
no telomerase activity was detected in cells prior to  
crisis when telomere length decreases, but abundant levels  
10 were detected in post-crisis cells which maintain telomere  
length. In contrast, the levels of *hTR* remained  
essentially constant throughout the immortalization  
process. The levels of *hEST2* RNA, on the other hand,  
parallel telomerase activity during cell immortalization.  
15 *hEST2* RNA was undetectable in the pre-crisis cells but was  
clearly present in the post-crisis, telomerase-positive  
cells. Induction of *hEST2* message is the rate-limiting  
step for the activation of telomerase during  
immortalization.

20 **EXAMPLE 7: Down-regulation of *hEST2* Expression upon  
Cellular Differentiation**

The up-regulation of *hEST2* RNA is associated with the  
activation of telomerase during cell immortalization. It  
remained unclear whether conditions that repress telomerase  
25 activity in cultured cells might similarly operate by  
shutting down expression of the *hEST2* mRNA in these cells.  
Human HL-60 promyelocytic leukemia cells can be induced to  
differentiate to mature granulocytes by treatment with  
retinoic acid. During this process, telomerase activity  
30 has been shown to decline over a period of two to five days  
(Sharma et al., *Proc. Natl. Acad. Sci., USA* 92:12343-12346

-65-

(1995); Bestilny et al., *Cancer Res.* 56:3796-3802 (1996); Savovsky et al., *Biochem. Biophys. Res. Commun.* 226:329-334 (1996); Xu et al., *Leukemia* 10:1354-1357 (1996)).

Induction of HL-60 differentiation was shown to lead  
5 to the disappearance of the *hEST2* mRNA within 24 hours,  
foreshadowing the loss of telomerase activity which  
declines more slowly. Once again, this contrasts with the  
behavior of the human telomerase RNA subunit, whose  
expression during induced differentiation remains constant,  
10 as also previously reported (Bestilny et al., *Cancer Res.*  
56:3796-3802 (1996)). Results presented here show that the  
levels of the *hEST2* mRNA decline precipitously within the  
first three hours after induced differentiation. This  
rapid down-modulation of is compatible with a short  
15 half-life for the *hEST2* message and suggests that the  
levels of this RNA species are under tight control in these  
cells. The contrasting delay in the decline of telomerase  
activity is consistent with the reported ~24 hour  
half-life of the enzymatic activity after cycloheximide  
20 treatment (Holt et al., *Mol. Cell. Biol.* 16:2932-2939  
(1996)).

#### EXAMPLE 8 Ectopic Expression of hTERT mRNA in Telomerase-Negative Cells and Assessment of Expression on Levels of Telomerase Activity

25 The *hTERT* cDNA was introduced into a mammalian  
expression construct carrying the CMV promoter. To  
distinguish the ectopically expressed hTERT protein from  
its endogenous counterpart, the C-terminus of the  
vector-encoded protein was marked with an influenza virus  
30 hemagglutinin (HA) epitope tag, yielding the plasmid

-66-

pCI-neo-*hTERT*-HA. Such a modification does not affect the catalytic activity of the Est2p protein of *S. cerevisiae* (Counter, C.M., et al., "The catalytic subunit of yeast telomerase", *Proc. Natl. Acad. Sci. USA*, 94:9202-9207  
5 (1997).

This *hTERT*-HA construct and a control empty vector were transfected into cells of the SV40-transformed GM847 human fibroblast line (Pereira-Smith, O.M., and Smith, J.R., "Genetic analysis of indefinite division in human  
10 cells: identification of four complementation groups", *Proc. Natl. Acad. Sci. USA*, 85:6042-6046 (1988). In contrast to most immortal human cells, which appear to activate telomerase in order to maintain telomere length (Shay, J.W., and Bacchetti, S., "A survey of telomerase  
15 activity in human cancer", *Eur. J. Cancer*, 33:787-791 (1997), the immortal GM847 cells are telomerase-negative (Bryan, T.M., et al., "Telomere elongation in immortal human cells without detectable telomerase activity", *EMBO J.*, 14:4240-4248 (1995). Moreover, while the *hTR* gene is  
20 transcribed in these cells (Bryan, T.M., et al., "The telomere lengthening mechanism in telomerase-negative immortal human cells does not involve the telomerase RNA subunit", *Hum. Mol. Genet.*, 6:921-926 (1997), they lack detectable levels of *hTERT* mRNA (Kilian, A., et al.,  
25 "Isolation of a candidate human telomerase catalytic subunit gene, which reveals complex splicing patterns in different cell types", *Hum. Mol. Genet.*, 6:2011-2019 (1997). These observations have led to the conclusion that GM847 cells employ a telomerase-independent mechanism to  
30 maintain telomere length (Murnane, J.P., et al., "Telomere dynamics in an immortal human cell line", *EMBO J.*,

-67-

13:4953-4962 (1994); (Bryan, T.M., et al., "Telomere elongation in immortal human cells without detectable telomerase activity", *EMBO J.*, 14:4240-4248 (1995), perhaps analogous to the recombination-based pathway used by yeast  
5 cells to allow chromosomes to maintain telomeres in the absence of a functional telomerase pathway (Lundblad, V., and Blackburn, E.H., "An alternative pathway for yeast telomere maintenance rescues est1- senescence. *Cell* 73, 347-360 (1993). Unlike normal, telomerase-negative human  
10 somatic cells, which lack replicative immortality, these GM847 cells could be propagated indefinitely following transfection, allowing us to study the properties of clonally isolated cell populations that have stably acquired the introduced *hTERT* gene.

15 A number of stably transfected GM847 cell clones were generated with either the control vector or the *hTERT*-HA expression vector, mRNA expression of *hTERT* was analyzed by RNase protection using probes that specifically recognize either the transfected *hTERT*-HA mRNA or the endogenous  
20 *hTERT* transcript. As expected, *hTERT*-HA transcript was detected only in GM847 sublines stably transfected with an *hTERT*-HA expression plasmid, but not in untransfected telomerase-positive control cell lines 293 and HL-60, in the parental GM847 line or in GM847 sublines transfected  
25 with the empty vector. The cells expressing *hTERT*-HA did not express the endogenous *hTERT* transcript, despite the fact that this RNA is clearly detected in telomerase-positive cells. Lastly, in accord with previous observations that the level of the *hTR* RNA subunit of  
30 telomerase does not correlate with enzyme activity, this RNA was detected in all cells tested, irrespective of whether the cells had telomerase activity. An actin

-68-

control probe demonstrates comparable loading of RNA from each cell line. Moreover, the specificity of the probes used was demonstrated by their failure to protect tRNA.

Expression of hTERT-HA was also analysed at the protein level, by immunoblotting with an anti-HA antibody probe directed against the HA tag of the vector-encoded hTERT protein. A ~ 130 kDa product corresponding to the predicted size of hTERT was detected in those lines derived from GM847 cells that were stably transfected with the hTERT-HA expression construct, but not in those cell clones that had been transfected with the empty vector. The antibody likewise did not detect endogenous (untagged) hTERT known to be expressed in 293 cells (Meyerson, M., et al. "hEST2, the Putative Human Telomerase Catalytic Subunit Gene, Is Up-Regulated in Tumor Cells and during Immortalization," *Cell*, 90:785-795 (1997), a telomerase-positive control cell line.

Assessment of whether these hitherto telomerase-negative GM847 cell lines acquired telomerase activity together with the stable ectopic expression of hTERT was carried out. Telomerase activity was measured in these different cell lines by assaying the ability of a cellular extract to elongate a primer in a telomerase-specific manner. The products of this *in vitro* reaction are subsequently detected by specific PCR amplification, yielding a ladder of products differing from one another by 6 bp (Kim, N.W., et al., "Specific association of human telomerase activity with immortal cells and cancer", *Science*, 266:2011-2015 (1994)). Telomerase activity was detected in 293 cells, a cell line with one of the highest levels of telomerase activity known. This activity was sensitive to heat treatment of the extract, which inactivates telomerase (Kim, N.W., et al., "Specific

-69-

association of human telomerase activity with immortal cells and cancer", *Science*, 266:2011-2015 (1994)). In contrast, almost no telomerase products were detectable following assay of extracts from untransfected GM847 cells or from GM847 sublines stably transfected with the empty control vector. This inability to detect telomerase was not due to the presence of a PCR-inhibiting activity, as an internal control was specifically PCR-amplified, (Kim, N.W., and Wu, F. "Advances in quantification and characterization of telomerase activity by the telomeric repeat amplification protocol (TRAP)", *Nucleic Acids Res.*, 25: 2595-2597 (1997)).

In marked contrast, telomerase activity was readily detectable in those clones of GM847 cells that were stably transfected with the *hTERT*-HA expression vector and this activity was sensitive to heat inactivation. The levels of telomerase activity observed in these transfectants approached those seen in extracts from 293 cells. The restoration of telomerase activity in the cells transfected with *hTERT*-HA was not due to the up-regulation of the endogenous *hTERT* gene, as evidenced by failure to detect the corresponding mRNA in these cells. Thus, ectopic expression of *hTERT* in previously telomerase-negative cells is sufficient to generate telomerase activity at levels comparable to those found in immortalized telomerase-positive cells.

The telomerase activity detected in cells transfected with the *hTERT*-HA expression vector was physically associated with ectopically produced *hTERT*-HA, confirming that *hTERT* is, as predicted, a constituent of the telomerase holoenzyme. Telomerase activity could be immunoprecipitated with an anti-HA monoclonal antibody from

-70-

extracts of either 293 cells or GM847 cells that ectopically express *hTERT*-HA. The telomerase activity was not immunoprecipitated with an antibody directed against an irrelevant antigen (anti-p53), nor was it

5 immunoprecipitated when either antibody was incubated with extracts prepared from control vector-transfected GM847 cells. Taken together, these data indicate that telomerase activity is specifically co-immunoprecipitated with *hTERT*-HA.

10 Whether the observed induction of telomerase activity following ectopic *hTERT* expression was unique to GM847 cells was also determined. This was done by transiently transfecting WI-38 normal human fibroblasts, which lack detectable levels of telomerase activity and *hTERT* message  
15 but express the *hTR* gene (Meyerson, M., et al., "*hEST2*, the Putative Human Telomerase Catalytic Subunit Gene, Is Up-Regulated in Tumor Cells and during Immortalization. Cell 90, 785-795 (1997), with the *hTERT*-HA expression construct. Since transient transfection of normal human  
20 cells is extremely inefficient, the plasmid pGreenLantern-1, which encodes the green fluorescent protein (GFP), was co-transfected with either the *hTERT*-HA expression vector or the control vector. Cells expressing GFP, and hence quite likely the co-transfected plasmid,  
25 were sorted by virtue of their fluorescence.

Extracts derived from both populations of fluorescing cells were assayed for telomerase activity. Whereas fibroblasts transfected with pGreenLantern-1 and pCI-neo vector alone lacked enzymatic activity, those  
30 co-transfected with the *hTERT*-HA expression vector were clearly telomerase-positive. Transfection with pCI-neo-*hTERT*-HA of IMR-90 cells, another telomerase-



-71-

negative normal human fibroblast cell strain that does not normally express *hTERT* (Kim, N.W., et al., "Specific association of human telomerase activity with immortal cells and cancer", *Science*, 266: 2011-2015 (1994);

- 5 (Meyerson, M., et al., "*hEST2*, the Putative Human Telomerase Catalytic Subunit Gene, Is Up-Regulated in Tumor Cells and during Immortalization", *Cell*, 90: 785-795 (1997), also gave rise to telomerase activity (not shown). Thus, ectopic expression of *hTERT* in these two types of normal
- 10 human somatic cells results in readily detectable telomerase activity.

Taken together, these data demonstrate that the ectopic expression of *hTERT* in otherwise telomerase-negative human cells is both necessary and

15 sufficient for induction of telomerase activity. Furthermore, the physical association of *hTERT* with telomerase activity confirms that *hTERT* is a telomerase subunit. The fact that forced expression of *hTERT* sufficed to impart telomerase activity indicates that levels of *hTR*

20 mRNA and TP-1 or other still unidentified components of the telomerase holoenzyme are not rate-limiting determinants of telomerase activity in these cells. Up-regulation of the *hTERT* gene is therefore the sole barrier to activation of telomerase in the tested cells.

- 25 It still remains to be determined if telomerase activity can be restored in this fashion in all telomerase-negative cells. However, results described here show that activity is conferred by ectopic expression of *hTERT* in cell types representative of two known classes of
- 30 telomerase-negative cells: telomerase-negative immortal cell lines and normal mortal human cell strains. Thus, it is likely that expression of *hTERT* mRNA is the

-72-

rate-determining step for telomerase activation in other human cells lacking enzyme activity. This makes the regulation of transcription from the *hTERT* promoter a potential target for modulation during tumorigenesis and cell immortalization.

#### Experimental Procedures

The following methods and materials were used in the experiments which are described above.

#### Radiation Hybrid Mapping

25 ng of genomic DNA from hybrid clones of the Genebridge 4 and Stanford G3 radiation hybrid (RH) mapping panels (Research Genetics, Inc. Huntsville, AL) was PCR-amplified with the primer pair M1 (forward-5' -CACAGCCAGGCCGAGAGCAGA- 3' (SEQ ID NO.: 48) and reverse-5' -AGGCCTCAGCCGGACACTCAG- 3') (SEQ ID NO.: 49), yielding a 170 bp fragment in the 3'-untranslated region, and with primer pair M2 (forward-5' -GAAGAAAACATTTCTGTCGTG- 3' (SEQ ID NO.: 50) and reverse- 5' -GCCCTTGGCCCCCAGCGACAT- 3') (SEQ ID NO.: 51), generating a 180 bp fragment crossing a putative intron-exon boundary near the *hEST2* stop codon. PCR was carried out for 35 cycles of 94°C for 0.5 min., 69°C for M1 or 65°C for M2 for 0.5 min. and 72°C for 1.5 min. The results of the PCR screening were analyzed using the statistical program RHMAP provided through the following two e-mail servers on the World Wide Web:  
<http://shgc-www.stanford.edu> and  
<http://www-genome.wi.mit.edu/cgi-bin/contig/rhmapper.pl>.  
Linkage of STS markers to a physical chromosomal map was achieved by accessing the servers:  
<http://www.cephb.fr/ceph-genethon-map.html> and  
<http://www-genome.wi.mit.edu>. Comparison to YACs

-73-

physically mapped by FISH was achieved with the server:  
ftp://ftp.cephb.fr/pub/ceph-genethon-map/FISH/29MAR95.DAT.

#### RNA Isolation from Cell Lines and Primary Tissues

Cell lines were obtained from the American Type

- 5 Culture Collection and grown under standard conditions.  
For HL-60 differentiation assays, cells were pelleted and  
resuspended in normal growth medium plus all-trans retinoic  
acid (Sigma) at a final concentration of 1  $\mu$ M.

- 10 All primary normal and tumor tissues were obtained  
from the Massachusetts General Hospital tumor bank.  
Tissues were processed in a tissue homogenizer. RNA  
samples were prepared in the RNA Stat-60 solution (Tel-Test  
"B", Friendswood, Texas) according to the manufacturer's  
protocol.

#### 15 Northern Hybridization

- Duplicate filters containing poly A(+)-selected mRNAs  
from various human tissues and cell lines (Multiple Tissue  
Northern Blots, Clontech, Palo Alto, CA), were incubated  
according to the manufacturer's instructions with two  
20 independent *hEST2* probes. One probe was derived by  
PCR-amplifying plasmid 712562 with primers HT-1 and HT-5 to  
generate the 377 bp fragment described above; the second,  
more 3' probe was generated from the 1064 bp *Stu* I fragment  
of the same plasmid. The duplicate blots were then  
25 rehybridized with a  $\beta$ -actin probe. Northern blots for  
detection of *hTR* were performed with total RNA and an  
*hTR*-specific probe (Feng et al., *Science* 269:1236-1241  
(1995)).

#### RNAse Protection Analysis

- 30 Radiolabelled RNA probes for RNAse protection analysis  
were synthesized using [ $\alpha$ -<sup>32</sup>P]UTP, T7 RNA polymerase and

- 74 -

the MAXIscrip kit (Ambion, Austin, TX). DNA templates for probe synthesis were created as follows. For the *hEST2* probe, the insert from plasmid clone 712562 was subcloned into pUHD 10-3 (a gift from M. Gossen), and template DNA was synthesized by PCR using a forward plasmid-specific primer and a reverse primer containing 18bp of *hEST2* complementary sequence (5'-TCTCTGCGGAAGTTCTG) (SEQ ID NO.: 52) and the T7 promoter sequence. For the  $\beta$ -actin probe, template DNA was synthesized by PCR on a human  $\beta$ -actin cDNA insert (Clontech). The *hEST2* and the  $\beta$ -actin control probes were hybridized in the same reaction tube. The *hTR* probe was synthesized directly from the linearized pGRN83 plasmid (Feng et al., *Science* 269:1236-1241 (1995)).

RNAse protection analysis was performed using the HybSpeed RPA kit (Ambion) according to the manufacturer's protocol. Briefly, sample RNA and radiolabelled RNA probes were coprecipitated in ethanol, resuspended in hybridization buffer, hybridized at 68° C, then digested with RNases A and T1. Samples were then re-precipitated and analyzed on a 6% denaturing polyacrylamide gel.

#### Telomerase assays

Telomerase repeat amplification protocol (TRAP) assays were performed as described (Kim et al., *Science* 266:2011-2015 (1994)), in some cases using the TRAPeze telomerase detection kit (Oncor, Gaithersburg, MD).

#### EQUIVALENTS

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the

-75-

invention as defined by the appended claims. Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described

5 specifically herein. Such equivalents are intended to be encompassed in the scope of the claims.

-76-

## CLAIMS

What is claimed is:

1. Isolated DNA encoding the catalytic subunit of a eukaryotic telomerase holoenzyme.
- 5 2. Isolated DNA of Claim 1 which encodes the catalytic subunit of a yeast telomerase holoenzyme.
3. Isolated DNA of Claim 2 wherein the DNA is selected from the group consisting of: DNA of SEQ ID NO.: 1; DNA which hybridizes to the complement of SEQ ID NO.:  
10 1; and DNA which encodes the amino acid sequence of SEQ ID NO.: 2.
4. Isolated DNA of Claim 1 which encodes the catalytic subunit of a human telomerase holoenzyme.
5. Isolated DNA of Claim 4 wherein the DNA is selected  
15 from the group consisting of: DNA comprising the nucleotide sequence of SEQ ID NO.: 35; DNA which encodes hEST2 protein having the amino acid sequence of SEQ ID NO.: 3; and DNA comprising a nucleotide sequence which hybridizes to the complement of SEQ ID  
20 NO.: 35.
6. Isolated DNA of Claim 4 which is an isolated gene which localizes to human chromosome subband 5p15.33.
7. Isolated DNA comprising DNA encoding a human telomerase catalytic subunit, wherein the DNA encodes  
25 an RNA transcript which is up-regulated in a human tumor.

-77-

8. Isolated DNA of Claim 7 which is an isolated human gene which localizes to human chromosome subband 5p15.33.
- 5 9. Isolated DNA of Claim 7 which comprises *hEST2* cDNA of SEQ ID NO.: 35 or DNA which hybridizes to the complement of *hEST2* cDNA under conditions of high stringency.
- 10 10. Isolated DNA of Claim 7 wherein the human tumor is selected from the group consisting of: breast tumors, ovarian tumors, colon tumors and testicular tumors.
11. Isolated DNA of Claim 7 which encodes a human telomerase catalytic subunit which comprises approximately 1132 amino acid residues.
- 15 12. Isolated DNA of Claim 8 wherein the RNA transcript is also up-regulated in a cancer cell line, a telomerase-positive tissue or both.
13. Isolated mRNA which encodes a human telomerase catalytic subunit, wherein expression of the mRNA is up-regulated in a human tumor.
- 20 14. Isolated mRNA of Claim 13 wherein the human tumor is selected from the group consisting of: breast tumors, ovarian tumors, colon tumors and testicular tumors.
15. Isolated mRNA of Claim 13 wherein the mRNA has the sequence of SEQ ID NO.: 36.

-78-

16. A nucleic acid probe comprising DNA unique to *hEST2* DNA, *EST2* DNA and p123 DNA.
17. The nucleic acid probe of Claim 16 comprising DNA which encodes the telomerase motif of *hEST2* protein.
- 5 18. The nucleic acid probe of Claim 17 comprising nucleotides which encode amino acid residues 556 to 565 of SEQ ID NO.: 3 or nucleotides which encode amino acid residues 1 to 50 of *hEST2* protein or a portion thereof.
- 10 19. A method of assessing cells for malignancy or an increased likelihood of progression to malignancy, comprising:
  - a) obtaining cells to be assessed from an individual;
  - 15 b) processing cells obtained in a) to render DNA and/or RNA in the cells available for annealing of the DNA and/or RNA with complementary poly- or oligonucleotides, thereby producing processed cells;
  - 20 c) combining processed cells with *hEST2* DNA or RNA under conditions appropriate for annealing of DNA and/or RNA with complementary poly- or oligonucleotides, thereby producing a combination,
  - 25 wherein if annealing occurs in the combination produced in c), it is indicative of activation of telomerase and malignancy or increased likelihood of progression to malignancy.
20. The method of Claim 21 wherein the poly- or  
30 oligonucleotides are DNA which encodes the telomerase



- 79 -

motif or amino acid residues 1 to 50 of SEQ ID NO.: 3  
or a portion thereof.

21. A method of diagnosing or aiding in the diagnosis of  
development of malignancy in an individual,  
5 comprising:
- a) obtaining cells from the individual;
  - b) processing the cells obtained in a) to render  
proteins in the cells available for binding with  
antibodies;
  - 10 c) combining the product of b) with antibodies which  
bind hEST2 protein; and
  - d) determining whether binding occurs between the  
antibodies which bind hEST2 protein and protein  
in the cell,
  - 15 wherein if binding occurs, it is indicative of the  
presence of hEST2 protein in the cells and of  
malignancy or an increased likelihood of development  
of malignancy in the individual.
22. The method of Claim 21 wherein the antibodies which  
20 bind hEST2 protein bind the telomerase motif of hEST2  
protein or bind amino acid residues 1 to 50 of SEQ ID  
NO.: 3 or a portion thereof.
23. A method of reducing expression of *hEST2* RNA and hEST2  
protein in cells of an individual, comprising  
25 administering to the individual a drug selected from  
the group consisting of: drugs which inhibit or bind  
*hEST2* RNA and prevent or reduce production of hEST2  
protein and drugs which inhibit hEST2 protein function  
or activity.

-80-

24. The method of Claim 23 wherein the drug is selected from the group consisting of: small organic molecules; enzymes which degrade hEST2 protein; enzyme inhibitors; HEST2 transcriptional regulators; antisense molecules and dominant negative forms of hEST2 protein.
25. A method of treating cancer in an individual, comprising administering to the individual a drug which inhibits or binds hEST2 RNA and prevents or reduces production of hEST2 protein or a drug which inhibits hEST2 protein function or activity, under conditions appropriate for the drug to enter cells of the individual, whereby the drug inhibits or binds hEST2 RNA and prevents or reduces production of hEST2 protein or inhibits hEST2 protein function or activity.
26. The method of Claim 25 wherein the drug is selected from the group consisting of: enzymes which degrade hEST2 protein; enzyme inhibitors; hEST2 transcriptional regulators; antisense molecules and dominant negative forms of hEST2 protein.
27. A method of increasing lifespan of cells, comprising:  
(a) introducing into the cells hEST2 DNA, under conditions appropriate for expression of hEST2 DNA, whereby hEST2 protein is expressed or  
(b) introducing hEST2 protein into the cells, whereby sufficient hEST2 protein is expressed or present in the cells to increase lifespan of the cells.

-81-

28. The method of Claim 27, wherein the cells are cells in culture.
29. A method of decreasing lifespan of cells in an individual by reducing hEST2 protein function or activity, comprising administering to the individual a drug selected from the group consisting of:
- (a) drugs which inhibit or bind *hEST2* RNA and prevent or reduce production of hEST2 protein;
  - (b) drugs which inhibit hEST2 protein function or activity; and
  - (c) drugs which inhibit or bind *hEST2* DNA and prevent or reduce production of *hEST2* RNA transcript, wherein the drug is administered to the individual under conditions appropriate for entry of the drug into cells in sufficient quantity to reduce hEST2 protein function or activity.
30. The method of Claim 29 wherein the drug is selected from the group consisting of: small organic molecules; enzymes which degrade hEST2 protein; enzyme inhibitors; HEST2 transcriptional regulators; antisense molecules and dominant negative forms of hEST2 protein.

1/14

ATGAAAATCTTATTCGAGTTCAATTCAGAACCAAGCTTGACATTTGATCTACAGACCAACAGTACTTACAAGAGAAATTTAAATATGTGGTCACTTCAATGGCC  
M K I L F E F I Q D K L D I D L Q T N S T Y K E N L K C G H F N G  
TCGATGAATCTAATCTACGTGTTTCGCACCTACCAATTCAGAAATAGCATTACCATGCCCTTCTGGTGAGTAAAGCCACAAAGCAGTCAATGATCA  
L D E I L T C F A L P N S R K I A L P C L P G D L S H K A V I D H  
CTGCATCATTTACCTGTTGACGGCGAATATACAAACACGACTACAACTTGGCTATAAATAGTGAATAGAGATGTCAACAATAGTCTTTTTTTC  
C I I Y L L T G E L Y N V L T F G Y K I A R N E D V N N S L F C  
CATTCGCAATGTTAAGCTTACGTTGAAAGCGCTGCTTGGAAATGTTCCACAGTTTGGTCGGTACATACGCTTCTGTTGATTATGATCAAT  
H S A N V N V T L L K G A A W K M F H S L V G T Y A F V D L L I N  
ATACAGTAATCAATTAATGGCGAGTTTTCATCTCAATCGTGGGTACAGATGTAAAGAACCTCATCTGCGGCCCAATGGTCCACGATCATCCTC  
Y T V I Q F N G Q F F T Q I V G N R C N E P H L P P K W V Q R S S  
ATCATCCGCAACTGCTGCGAATCAACAACTTACAGAACCGAGTGACAAATAAACAATCTTACACAGCTCAATCAATAAATCTCTCTCTTTTCTCT  
S S A T A A Q I K Q L T E P V T N K Q F L H K L N I N S S F P  
TATAGCAAGTCTCTTCATCATCTATCAATAAGCTAACTGACTTGAGAGAGCTATTTTCCCAAAATTTGGTAAATTCCTCAGAGACTAA  
Y S K I L P S S S I K K L T D L R E A I F P T N L V K I P Q R L  
AGGTACGAATTAATTTGACGCTGCAAAAGCTATTAAAGAGACATAAGCGTTTGAAATPACGTTCTATTTTGAATAGTATTTGCCACCATTGGAGGGAC  
K V R I N L T L Q K L L K R H K R L N Y V S I L N S I C P P L E G T  
CGTATGGACTTGTCGCAATTTGAGTAGGCAATCACCAGGAGACGAGTCTTGAATTTACCTTAAATGGCTATTTTACAGAAATTTACCCCAAGAAATGTTT  
V L D L S H L S R Q S P K E R V L K F I I V I L Q K L L P Q E M F  
GGCTCAAGGAAATTTTCGGTGGTTGTTTCATTTCTGATATTTGGTTACCAAGCACAAATTTGAACACTTGAATCAATTTGGCGATTTGTTTCATTTCTCTG  
TAAGATTAAAGGATTTTCGGTGGTTGTTTCATTTCTGATATTTGGTTACCAAGCACAAATTTGAACACTTGAATCAATTTGGCGATTTGTTTCATTTCTCTG  
L R L K D F R W L F I S D I W F T K H N F E N L N Q L A I C F I S W  
GCTATTAGACAACTAATCCCAAAATATACAGACTTTTTTTTACTGACCGAATATCTTCTACAGTGACAAATTTGTTTACTTTAGACATGATCTTGG  
L F R Q L I P K I I Q T F F Y C T E I S S T V T I V Y F R H D T W  
AATAAATTTATCCCTTTTATCGTAGAATATTTTAAAGACGTACTTAGTTCGAAACCAACGATAGTGAACCAATATAGTTACAGTTGTCCCAATTTCA  
N K L I T P F I V E Y F K T Y L V E N N V C R N H N S Y T L S N F  
ATCATAGCAAAATGAGGATTATACCAAAAAAGTAATAATGAGTTTCAGGATTTATGCCATCCCATGCGAGGGGCGAGCGAAGAGAAATTCACAATTTA  
N H S K M R I I P K S N N E F R I I A I P C R G A D E E E F T I Y  
TAAGAGAAATCACAAAATGCTATCCAGCCCACTCAAAAATTTTAGAATACCTAAGAAACAAAGGCGGACCTAGTTTACTAAATATATTTCTCCAACG  
K E N H K N A I Q P T Q K I L E Y L R N K R P T S F T K I Y S P T  
CAATAGCTGACCGGTATCAAGAATTTAAGCAGAGACTTTTAAAGAAATTTAATAATGTCTTACCAGAGCTTTATTTATGAAATTTGATGTCAATCTT  
Q I A D R I K E F K Q R L L K K F N N V L P E L Y F M K F D V K S  
GCTATGATTCATACCAAGGATGGAATGTATGAGGATACCTCAAGGATGCGCTAAATAATGAAATGGTTCGTTAGATCTCAATATTTCTCAATAC  
C Y D S I P R M E C M R I L K D A L K N E N G F F V R S Q Y F F N T  
CAATACAGGTGTTAAGTTATTTAATGTGTAACTAGCAGAGTACCAAACTTATGAGCTATAGATAGATAATGTGAGGACGGTTCATTTATCA  
N T G V L K L F N V N A S R V P K P Y E L Y I D N V R T V H L S  
AATCAGGATGTTATAACGTTGTAGAGATGGAATATTTAAACAGCTTTGTGGTTGAAGTAAAGTGTACATAGAGAGATGGTCTTTTTCAGGGCT  
N Q D V I N V V E M E I F K T A L W V E D K C Y I R E D G L F Q G

FIG. 1A

2/14

CTAGTTTATCTGCTCCGATCGTTGATTTGGTGTATGACGATCTTCTGGAGTTTATAGCGAGTTTAAAGCCAGTCTTAGCCAGGACACATTAATTTTAA  
S L S A P I V D L V Y D D L L E F Y S E F K A S P S Q D T L I L K  
ACTGGCTGACGATTTCCCTTATATATCAACAGACCAACAGCAAGTGAATCAATATCAAAAGCTTGCCATGGCGGATTTCAAAAATATAATGCGAAAGCC  
L A D D F L I I S T D Q Q Q V I N I K K L A M G G F Q K Y N A K A  
AATAGAGACAAAATTTTAGCCGTAAGCTCCCAATCAGATGATGATACGGTTATTCAATTTTGTGCAATGCACATATTTGTTAAAGAAATGGAAGTTTGGGA  
N R D K I L A V S S Q S D D D T V I Q F C A M H I F V K E L E V W  
AACATTCAAGCACAAATGAATAATTTCCATATCCGTTTCAAAATCTAGTAAAGGATATTTGCAAGTTTAAATAGCGCTGTTTAAACACTAGATCTCTTATAA  
K H S S T M N N F H I R S K S K G I F R S L I A L F N T R I S Y K  
AACAAATTGACACAAAATTTAAATTCACAAACACCGTTCTCATGCAATTTGATCATGTTGTAAAGAACATTTCCGGAATGTTTAAATCTGCTTTTAAAGGAT  
T I D T N L N S T N T V L M Q I D H V V K N I S E C Y K S A F K D  
CTATCAATTAATGTTACGCAAAATATGCAATTTTCATTCTTTACACGCATCATTTGAATGACAGTCAGCGGTTGTCCAATTTACGAATGATGATCCTT  
L S I N V T Q N M Q F H S F L Q R I I E M T V S G C P I T K C D P  
TAATCGAGTATGAGGTACGATTCACCATATTTGAATGGATTTTGGAAAGCCTATCTTCAACACACATCAAAATTTAAAGATAATATCATTTCTTTTGAGAAA  
L I E Y E V R F T I L N G F L E S L S N T S K F K D N I I L L R K  
GGAAATTCAACTTGCAAGCATATATATATATATACATATAGTTAATTAG 2655  
E I Q H L Q A Y I Y I I H I I V N

FIG. 1B

3/14

	I	M P R A P R C R A V R S L L R S H Y R E V L P L A T F V R R L G P Q G W R L V Q R G D P A A F R A L H E S T 2
551		V A Q C L V C V P W D A R P P A A P S F R Q V S C L K E I E V A R V L Q R L C E I E G A K N V L A F G F A L D G A R G G H E S T 2
11		M E V D V D N Q A D N H G I H S A L K T C E E I K E A K T I Y S - W I Q K V I - R C R N Q S Q S H Y K D I E D I K I F A P 123
11		- - - - - M K I I F E - F I Q D K L D I D L Q T N S T Y K E N K C G H F N G E S t 2 p
1111		P P E A F T T S V S V L P N T V T D A L R G S G A W L G L L R R - V G O D V C V H E L A R C A L F V L V A P S C A Y O H E S T 2
939		Q T N V A P - P D H N E E D F K V I A R K E V F T G M I E - L I D K V L E I L S S - - - - - S D V S D R Q K L O P 123
584		L D E I L I T - - - - C F A L P N S R K I A L P C L P G D L S H K A V I D H G I I Y L L - - - - - T G E L Y L N E S t 2 p
170		N C G P P L Y O G A A T Q R P - P P H J S G P R R R L G C E R A W N H S V R E A G V P L G L P P G A R R R G G S A H E S T 2
1113		C F G - - - - F Q L R G N Q L A K T H L L T A L S T Q K Y F F Q D E W N - Q V R - - - - - A M I G N E L F R H L P 123
181		N L T - - - - - F S G Y K I A R - - - - - N E D V N N S L - - - - - - - - - F E s t 2 p
229		S R S L P L P K P P R R R G A P E P E R T P V C Q G S W A P G R T R G P S D R G F C V V S P A R P A E E T S L G A H E S T 2
160		Y T K Y L I F Q R T S E G T L V Q - - - - F C G N N V D H K V N D K F D K K - - - - - Q K G G A D M N E R P 123
100		C H S A N V N V T L L K G A W K M F H S L V G T Y A F V D L I N Y T V I Q E N - - - - - G Q F F T I Q V G N R C N E P H E S t 2 p
289		D S G T R H S H P S V G R Q A H A G P P S T S R P M P D T C P P Y A E T K H L Y S S G D K - E Q L R P S F L L H E S T 2
208		C C S T C K Y N V K N E K D H F - - - - L N N I N V P N - W N - - - - - M K S R T R I E Y C T H F N R N N Q F K K H E F P 123
1157		L P P K W V Q R S S S S A T A A - Q I K Q L T E P - V - - - - - T N K Q F L H K L N I N S S F F E P Y S K I E s t 2 p
348		S S L R P S L T G A R R L V E T I F L G S R P W M P G T P R L P R L P Q R Y W Q M R P L F E L L G N H A Q C P Y I G V H E S T 2
260		V S N K N N I S A M D R - A Q T I F - - - - T N I F R F N I R K - K L K D K V I E K I A Y M E K V K D F - - - - - N F N Y P 123
205		L P S S S S I K K L D T D R E A T F - P T N L V K I P Q - - - - - R L K V R I N T L Q K L K R H K R L N - - - - - X V S E s t 2 p
408		L U K T H C P P R A A V T P A A G V C A R E K P Q G S V A A P E E E D T D P R R L V Q L R Q H S S P W Q V Y G - F V R H E S T 2
312		Y H T K S C P L - - - - - P E N W R C K Q I E N L I N K T R E E K S - - - - - K Y Y E E I F S Y T D N K C V T Q P I N P 123
256		I U N S I D P - - - - - P L E G T V L D - - - - - L S H L S R Q S P K E R V L - K F I I E s t 2 p
467		A C R R R V P P G L W G S R H N E R R L R N T K K F I S L G K H A K L S L Q E L T W K M S V R G C A W L R R S P G V H E S T 2
364		E F F Y N I P K D F L T G R - W R K N F Q K K V K K Y V E L N K H E L I H K N L L E K I N T R E I S W M Q V E T S A P 123
289		V I L Q K L P Q E M F G S K K K N K G K I I N L N L L S L P L N G Y L P F D S L K K L R L K D F R W L F I - - - - - S D E s t 2 p
527		G C V P A A E H R L R E E I A K F L H W L M S V V V E L R I S F Y V T E I T F Q K N R L F F Y R K S V S W S K L Q S H E S T 2
343		K V Y F A E D H E - N I Y V W K L E R W I E D V V S L T R I C F I F Q Q K S Y S K T Y N Y R K N I W D I M K P 123
427		I W T K H F N F N L N Q A I C F I S W L F R Q L I P K I I Q T F E Y C I T E I S S S T - T I V F R H N I D T W N L I T E S t 2 p

**FIG. 2A**

4/14

587 I G R Q H R V Q L S L R R Q H R E A R -- P A L L T S R R P I P K -- P D G L R P I - V N M D Y - V V H E S T 2  
 588 M S R - A D C K E T A E V Q E E E E W K K S -- -- L G A P G K L R L I P K -- K T T F R P I - M T F N - -- p123  
 589 P F R - V E Y F K T Y V -- -- E N N V C R N H N S Y T L S N F H S K M R I L P K K S N N E E R I L A I P C R G A D E E s t 2 p  
 590  
 591 G A R F R R R K K R A E R L S R V K A T F V N Y E R A R R P G L -- -- L G A S V L G L D D L H R A W R T L L R V H E S T 2  
 592 K - K I V N S O R K T T K L E T N T L L N S H M L K T L K N R M F K D P F G F A V F N Y D D V M K K Y E E F V C K W p123  
 593 E E F I Y K E N H K N A I Q P T Q K I L - E Y R N K R P T S F T - K I Y S P T Q I A - D R I K E F K Q R L L K K F E s t 2 p  
 594  
 595  
 596  
 597  
 598 R A G D P P P E L Y F V K V D V T G A Y D I L P Q D R L T E V I -- -- -- -- -- K S I I R P Q N T Y C V R R H E S T 2  
 599 K - V V G Q P K L F F A T M D L E K Y D S V N R E K G S T F F K T T K L L S S D F W I M T A Q I K R K N N I V I D S p123  
 600 -- -- N N V L P E L Y F E M R F D V K S C Y D S I P R M E C R I L K -- -- -- -- -- D A L K N E G F F V R S E s t 2 p  
 601  
 602  
 603  
 604  
 605  
 606  
 607  
 608  
 609  
 610  
 611  
 612  
 613  
 614  
 615  
 616  
 617  
 618  
 619  
 620  
 621  
 622  
 623  
 624  
 625  
 626  
 627  
 628  
 629  
 630  
 631  
 632  
 633  
 634  
 635  
 636  
 637  
 638  
 639  
 640  
 641  
 642  
 643  
 644  
 645  
 646  
 647  
 648  
 649  
 650  
 651  
 652  
 653  
 654  
 655  
 656  
 657  
 658  
 659  
 660  
 661  
 662  
 663  
 664  
 665  
 666  
 667  
 668  
 669  
 670  
 671  
 672  
 673  
 674  
 675  
 676  
 677  
 678  
 679  
 680  
 681  
 682  
 683  
 684  
 685  
 686  
 687  
 688  
 689  
 690  
 691  
 692  
 693  
 694  
 695  
 696  
 697  
 698  
 699  
 700  
 701  
 702  
 703  
 704  
 705  
 706  
 707  
 708  
 709  
 710  
 711  
 712  
 713  
 714  
 715  
 716  
 717  
 718  
 719  
 720  
 721  
 722  
 723  
 724  
 725  
 726  
 727  
 728  
 729  
 730  
 731  
 732  
 733  
 734  
 735  
 736  
 737  
 738  
 739  
 740  
 741  
 742  
 743  
 744  
 745  
 746  
 747  
 748  
 749  
 750  
 751  
 752  
 753  
 754  
 755  
 756  
 757  
 758  
 759  
 760  
 761  
 762  
 763  
 764  
 765  
 766  
 767  
 768  
 769  
 770  
 771  
 772  
 773  
 774  
 775  
 776  
 777  
 778  
 779  
 780  
 781  
 782  
 783  
 784  
 785  
 786  
 787  
 788  
 789  
 790  
 791  
 792  
 793  
 794  
 795  
 796  
 797  
 798  
 799  
 800  
 801  
 802  
 803  
 804  
 805  
 806  
 807  
 808  
 809  
 810  
 811  
 812  
 813  
 814  
 815  
 816  
 817  
 818  
 819  
 820  
 821  
 822  
 823  
 824  
 825  
 826  
 827  
 828  
 829  
 830  
 831  
 832  
 833  
 834  
 835  
 836  
 837  
 838  
 839  
 840  
 841  
 842  
 843  
 844  
 845  
 846  
 847  
 848  
 849  
 850  
 851  
 852  
 853  
 854  
 855  
 856  
 857  
 858  
 859  
 860  
 861  
 862  
 863  
 864  
 865  
 866  
 867  
 868  
 869  
 870  
 871  
 872  
 873  
 874  
 875  
 876  
 877  
 878  
 879  
 880  
 881  
 882  
 883  
 884  
 885  
 886  
 887  
 888  
 889  
 890  
 891  
 892  
 893  
 894  
 895  
 896  
 897  
 898  
 899  
 900  
 901  
 902  
 903  
 904  
 905  
 906  
 907  
 908  
 909  
 910  
 911  
 912  
 913  
 914  
 915  
 916  
 917  
 918  
 919  
 920  
 921  
 922  
 923  
 924  
 925  
 926  
 927  
 928  
 929  
 930  
 931  
 932  
 933  
 934  
 935  
 936  
 937  
 938  
 939  
 940  
 941  
 942  
 943  
 944  
 945  
 946  
 947  
 948  
 949  
 950  
 951  
 952  
 953  
 954  
 955  
 956  
 957  
 958  
 959  
 960  
 961  
 962  
 963  
 964  
 965  
 966  
 967  
 968  
 969  
 970  
 971  
 972  
 973  
 974  
 975  
 976  
 977  
 978  
 979  
 980  
 981  
 982  
 983  
 984  
 985  
 986  
 987  
 988  
 989  
 990  
 991  
 992  
 993  
 994  
 995  
 996  
 997  
 998  
 999  
 1000  
 1001  
 1002  
 1003  
 1004  
 1005  
 1006  
 1007  
 1008  
 1009  
 1010  
 1011  
 1012  
 1013  
 1014  
 1015  
 1016  
 1017  
 1018  
 1019  
 1020  
 1021  
 1022  
 1023  
 1024  
 1025  
 1026  
 1027  
 1028  
 1029  
 1030  
 1031  
 1032  
 1033  
 1034  
 1035  
 1036  
 1037  
 1038  
 1039  
 1040  
 1041  
 1042  
 1043  
 1044  
 1045  
 1046  
 1047  
 1048  
 1049  
 1050  
 1051  
 1052  
 1053  
 1054  
 1055  
 1056  
 1057  
 1058  
 1059  
 1060  
 1061  
 1062  
 1063  
 1064  
 1065  
 1066  
 1067  
 1068  
 1069  
 1070  
 1071  
 1072  
 1073  
 1074  
 1075  
 1076  
 1077  
 1078  
 1079  
 1080  
 1081  
 1082  
 1083  
 1084  
 1085  
 1086  
 1087  
 1088  
 1089  
 1090  
 1091  
 1092  
 1093  
 1094  
 1095  
 1096  
 1097  
 1098  
 1099  
 1100  
 1101  
 1102  
 1103  
 1104  
 1105  
 1106  
 1107  
 1108  
 1109  
 1110  
 1111  
 1112  
 1113  
 1114  
 1115  
 1116  
 1117  
 1118  
 1119  
 1120  
 1121  
 1122  
 1123  
 1124  
 1125  
 1126  
 1127  
 1128  
 1129  
 1130  
 1131  
 1132  
 1133  
 1134  
 1135  
 1136  
 1137  
 1138  
 1139  
 1140  
 1141  
 1142  
 1143  
 1144  
 1145  
 1146  
 1147  
 1148  
 1149  
 1150  
 1151  
 1152  
 1153  
 1154  
 1155  
 1156  
 1157  
 1158  
 1159  
 1160  
 1161  
 1162  
 1163  
 1164  
 1165  
 1166  
 1167  
 1168  
 1169  
 1170  
 1171  
 1172  
 1173  
 1174  
 1175  
 1176  
 1177  
 1178  
 1179  
 1180  
 1181  
 1182  
 1183  
 1184  
 1185  
 1186  
 1187  
 1188  
 1189  
 1190  
 1191  
 1192  
 1193  
 1194  
 1195  
 1196  
 1197  
 1198  
 1199  
 1200  
 1201  
 1202  
 1203  
 1204  
 1205  
 1206  
 1207  
 1208  
 1209  
 1210  
 1211  
 1212  
 1213  
 1214  
 1215  
 1216  
 1217  
 1218  
 1219  
 1220  
 1221  
 1222  
 1223  
 1224  
 1225  
 1226  
 1227  
 1228  
 1229  
 1230  
 1231  
 1232  
 1233  
 1234  
 1235  
 1236  
 1237  
 1238  
 1239  
 1240  
 1241  
 1242  
 1243  
 1244  
 1245  
 1246  
 1247  
 1248  
 1249  
 1250  
 1251  
 1252  
 1253  
 1254  
 1255  
 1256  
 1257  
 1258  
 1259  
 1260  
 1261  
 1262  
 1263  
 1264  
 1265  
 1266  
 1267  
 1268  
 1269  
 1270  
 1271  
 1272  
 1273  
 1274  
 1275  
 1276  
 1277  
 1278  
 1279  
 1280  
 1281  
 1282  
 1283  
 1284  
 1285  
 1286  
 1287  
 1288  
 1289  
 1290  
 1291  
 1292  
 1293  
 1294  
 1295  
 1296  
 1297  
 1298  
 1299  
 1300  
 1301  
 1302  
 1303  
 1304  
 1305  
 1306  
 1307  
 1308  
 1309  
 1310  
 1311  
 1312  
 1313  
 1314  
 1315  
 1316  
 1317  
 1318  
 1319  
 1320  
 1321  
 1322  
 1323  
 1324  
 1325  
 1326  
 1327  
 1328  
 1329  
 1330  
 1331  
 1332  
 1333  
 1334  
 1335  
 1336  
 1337  
 1338  
 1339  
 1340  
 1341  
 1342  
 1343  
 1344  
 1345  
 1346  
 1347  
 1348  
 1349  
 1350  
 1351  
 1352  
 1353  
 1354  
 1355  
 1356  
 1357  
 1358  
 1359  
 1360  
 1361  
 1362  
 1363  
 1364  
 1365  
 1366  
 1367  
 1368  
 1369  
 1370  
 1371  
 1372  
 1373  
 1374  
 1375  
 1376  
 1377  
 1378  
 1379  
 1380  
 1381  
 1382  
 1383  
 1384  
 1385  
 1386  
 1387  
 1388  
 1389  
 1390  
 1391  
 1392  
 1393  
 1394  
 1395  
 1396  
 1397  
 1398  
 1399  
 1400  
 1401  
 1402  
 1403  
 1404  
 1405  
 1406  
 1407  
 1408  
 1409  
 1410  
 1411  
 1412  
 1413  
 1414  
 1415  
 1416  
 1417  
 1418  
 1419  
 1420  
 1421  
 1422  
 1423  
 1424  
 1425  
 1426  
 1427  
 1428  
 1429  
 1430  
 1431  
 1432  
 1433  
 1434  
 1435  
 1436  
 1437  
 1438  
 1439  
 1440  
 1441  
 1442  
 1443  
 1444  
 1445  
 1446  
 1447  
 1448  
 1449  
 1450  
 1451  
 1452  
 1453  
 1454  
 1455  
 1456  
 1457  
 1458  
 1459  
 1460  
 1461  
 1462  
 1463  
 1464  
 1465  
 1466  
 1467  
 1468  
 1469  
 1470  
 1471  
 1472  
 1473  
 1474  
 1475  
 1476  
 1477  
 1478  
 1479  
 1480  
 1481  
 1482  
 1483  
 1484  
 1485  
 1486  
 1487  
 1488  
 1489  
 1490  
 1491  
 1492  
 1493  
 1494  
 1495  
 1496  
 1497  
 1498  
 1499  
 1500  
 1501  
 1502  
 1503  
 1504  
 1505  
 1506  
 1507  
 1508  
 1509  
 1510  
 1511  
 1512  
 1513  
 1514  
 1515  
 1516  
 1517  
 1518  
 1519  
 1520  
 1521  
 1522  
 1523  
 1524  
 1525  
 1526  
 1527  
 1528  
 1529  
 1530  
 1531  
 1532  
 1533  
 1534  
 1535  
 1536  
 1537  
 1538  
 1539  
 1540  
 1541  
 1542  
 1543  
 1544  
 1545  
 1546  
 1547  
 1548  
 1549  
 1550  
 1551  
 1552  
 1553  
 1554  
 1555  
 1556  
 1557  
 1558  
 1559  
 1560  
 1561  
 1562  
 1563  
 1564  
 1565  
 1566  
 1567  
 1568  
 1569  
 1570  
 1571  
 1572  
 1573  
 1574  
 1575  
 1576  
 1577  
 1578  
 1579  
 1580  
 1581  
 1582  
 1583  
 1584  
 1585  
 1586  
 1587  
 1588  
 1589  
 1590  
 1591  
 1592  
 1593  
 1594  
 1595  
 1596  
 1597  
 1598  
 1599  
 1600  
 1601  
 1602  
 1603  
 1604  
 1605  
 1606  
 1607  
 1608  
 1609  
 1610  
 1611  
 1612  
 1613  
 1614  
 1615  
 1616  
 1617  
 1618  
 1619  
 1620  
 1621  
 1622  
 1623  
 1624  
 1625  
 1626  
 1627  
 1628  
 1629  
 1630  
 1631  
 1632  
 1633  
 1634  
 1635  
 1636  
 1637  
 1638  
 1639  
 1640  
 1641  
 1642  
 1643  
 1644  
 1645  
 1646  
 1647  
 1648  
 1649  
 1650  
 1651  
 1652  
 1653  
 1654  
 1655  
 1656  
 1657  
 1658  
 1659  
 1660  
 1661  
 1662  
 1663  
 1664  
 1665  
 1666  
 1667  
 1668  
 1669  
 1670  
 1671  
 1672  
 1673  
 1674  
 1675  
 1676  
 1677  
 1678  
 1679  
 1680  
 1681  
 1682  
 1683  
 1684  
 1685  
 1686  
 1687  
 1688  
 1689  
 1690  
 1691  
 1692  
 1693  
 1694  
 1695  
 1696  
 1697  
 1698  
 1699  
 1700  
 1701  
 1702  
 1703  
 1704  
 1705  
 1706  
 1707  
 1708  
 1709  
 1710  
 1711  
 1712  
 1713  
 1714  
 1715  
 1716  
 1717  
 1718  
 1719  
 1720  
 1721  
 1722  
 1723  
 1724  
 1725  
 1726  
 1727  
 1728  
 1729  
 1730  
 1731  
 1732  
 1733  
 1734  
 1735  
 1736  
 1737  
 1738  
 1739  
 1740  
 1741  
 1742  
 1743  
 1744  
 1745  
 1746  
 1747  
 1748  
 1749  
 1750  
 1751  
 1752  
 1753  
 1754  
 1755  
 1756  
 1757  
 1758  
 1759  
 1760  
 1761  
 1762  
 1763  
 1764  
 1765  
 1766  
 1767  
 1768  
 1769  
 1770  
 1771  
 1772  
 1773  
 1774  
 1775  
 1776  
 1777  
 1778  
 1779  
 1780  
 1781  
 1782  
 1783  
 1784  
 1785  
 1786  
 1787  
 1788  
 1789  
 1790  
 1791  
 1792  
 1793  
 1794  
 1795  
 1796  
 1797  
 1798  
 1799  
 1800  
 1801  
 1802  
 1803  
 1804  
 1805  
 1806  
 1807  
 1808  
 1809  
 1810  
 1811  
 1812  
 1813  
 1814  
 1815  
 1816  
 1817  
 1818  
 1819  
 1820  
 1821  
 1822  
 1823  
 1824  
 1825  
 1826  
 1827  
 1828  
 1829  
 1830  
 1831  
 1832  
 1833  
 1834  
 1835  
 1836  
 1837  
 1838  
 1839  
 1840  
 1841  
 1842  
 1843  
 1844  
 1845  
 1846  
 1847  
 1848  
 1849  
 1850  
 1851  
 1852  
 1853  
 1854  
 1855  
 1856  
 1857  
 1858  
 1859  
 1860  
 1861  
 1862  
 1863  
 1864  
 1865  
 1866  
 1867  
 1868  
 1869  
 1870  
 1871  
 1872  
 1873  
 1874  
 1875  
 1876  
 1877  
 1878  
 1879  
 1880  
 1881  
 1882  
 1883  
 1884  
 1885  
 1886  
 1887  
 1888  
 1889  
 1890  
 1891  
 1892  
 1893  
 1894  
 1895  
 1896  
 1897  
 1898  
 1899  
 1900  
 1901  
 1902  
 1903  
 1904  
 1905  
 1906  
 1907  
 1908  
 1909  
 1910  
 1911  
 1912  
 1913  
 1914  
 1915  
 1916  
 1917  
 1918  
 1919  
 1920  
 1921  
 1922  
 1923  
 1924  
 1925  
 1926  
 1927  
 1928  
 1929  
 1930  
 1931  
 1932  
 1

5/14




FIG. 3A

FIG. 3B

FIG. 3C





FIG. 3D

FIG. 3E

FIG. 3F



FIG. 4



10	20	30	40	50	60	70	80
CAGGCAGCGCTGCGTCTGCTGCGCACGCTGGGAAGCCCTGGCCCGCCGACCCCGCGGATGCGCGGCTGCCCGCTGCC	80						
GAGCGGTGCGTCTGCTGCGCACGCTACCGGAGGAGTGTGCTCCGCTGAGCAGCTGCTGGCGGCGCTGGGCCCCCAG	160						
CGCACGCGCGCCCGCCCGCCCTCTTCGCGCAGGTGTCTGCTGCTGAAGAGACTGGTGGCCCGGAGTGTGACAGGC	320						
TGTGCGAGCGCGGCGCGAAGAACGTGTGCGCTTCGCGCTCGCGCTGCTGAGACGGGGCCCCGCGGGSCCCCCCGAGGCC	400						
410	420	430	440	450	460	470	480
TTACACACCGCTGCGCAGCTACCTGTCCCAACACGCTGACCGACGCACTGCGGGGGAGCGGGCGTGGGGCTGCTGTT	480						
GCGCGCGTGGGACGACGTGTGTTTCACTGTGTGGACAGCTGGCGCTCTTTGTGCTGGTCCAGTCCGCTT	560						
ACCAGGTGTGCGGCGCGCTGTACCAAGCTCGGCGCTGCCACTCAGGCCCGGCCCGCCACACGCTAGTGGACCCCGA	640						
AGGCGTCTGGATGCGAACGGGCCCTGGAACCATAGCGTCAAGGAGGCGGGGTCCCCCTGGGCTTGCCAGCCCGGGTGC	720						
GAGGAGCGCGGGGCGAGTGCCACCGCAAGTCTGCGCTTGCCCAAGAGGCCCGAGCGCTGGCGTGTCCCCCTGAGCGCGAGC	800						
810	820	830	840	850	860	870	880
GGACGCCCGTTGGCGAGGGTCTTGGGCCACCCCGGACGACGCGTGGACCGAGTGACCTGCTGGTTCCTGTGTGGTGTCA	880						
CCTGCACACCCCGCGAAGAACCTCTTTGGAGGGTGGCTCTCTGBCACGCGCACTCCACCCACTCCGTGGGCGG	960						
CCAGCACACCGGGCCCCCATACACTCGCGGCCACACAGCTCCTGGGACACGCTTGTCCCCGGTGTACGCGGAGA	1040						
CCAAGCACTTCTCTACTCTCAGGCGACAAGGAGAGCTGGGCGCTCTCTTCTACTAGCTCTCTGAGGCGCCAGCCTG	1120						
ACTGGCGCTCGGAGGCTCGTGGAGACCATCTTTCTGGGTTCCAGGCCCTGGATGTCAGGGACTCCCCGCAAGGTTGCCCGG	1200						
1210	1220	1230	1240	1250	1260	1270	1280
CCTGCCCCAGCGCTACTGGCAATGTCGGGCCCTCTGTTCTGAGAGTGTCTGGGAACCAACGCGAGTGCCCTACGGGGTGC	1280						
TCTCAAGACGCACTGCCCGTGCAGCTGCGGTACCCACGACAGCCGGTGTCTGTGCCGAGAGAAGCCCCAGGGCTCT	1360						
GTGCGGCCCGCGAGGAGGAGACACAGACCCCGTGCCTGTGTGCTGCTGCGCACGACAGCAGCCCCCTGGCAGGT	1440						
GTCAGGCTTGTGCGGGCTGCTGCGCGGCTGGTGTGCCCCCAGGCTCTGGGGCTCCAGGCAACACGAACGCCGCTTCC	1520						
TCAGGAACACCAAGAAGTTCACTCTCCTGGGGAAGCATGTCCAAAGCTCTCGTGCAGGAGTGACCTGGAAGATGAGCGTG	1600						
1610	1620	1630	1640	1650	1660	1670	1680
CGGGCTGCGCTGGCTGGCTGCGAGGCCAGGGGTGGCTGTGTTTCGGCCGCGACAGACCCGTCTGCGTGAAGAGATCCT	1680						
GGCCAACTTCTGCTGCACTGGCTGATGAGTGTACGTGCTGAGCTGTCTCAGTCTTCTTTTATGTACGGAGACCCAGT	1760						
TTCAAAAGAACAGGCTCTTTTCTACCGGAAGAGTGTCTGGAGCAAGTTGCAAGCATTTGGAATCAGACAGCACTTGAAG	1840						
AGGTTGCACTGCGGAGCTGTGCGAAGACAGAGGTCAGGCAGCATCGGAAGCCAGGCCCGCTGTGACGTCCAGACT	1920						
CCGCTTTCATCCCCAAGCCTGACGGGCTGCGGCCGATTTGTGAACAATGGACTACGTCTGTGGGAGCCAGAACGTTTCCGCAGAG	2000						
2010	2020	2030	2040	2050	2060	2070	2080
AAAAGAGCGCAGGCTCTACCTCGAGGGTGAAGGCACTGTTACGCGTGTCTCAACTACGAGCGGGCGGCGCCCCCGG	2080						
CTCCTGGGCGCTGTGTGGCCTGGACGATATCCACAGGGCCTGGCGCACCTTCGTGTGCTGCTGTGCGGCCAGGA	2160						
CCGCGCGCTGAGCTGTACTTTGTCAAGAGTGGATGTGAACGGGCGGTGACGACCATCCCCAGGACAGGCTACGCGAGG	2240						
TATCGCGCAGCATCAAAACCCCAAGACAGCTACTGCTGTCGTCGTTGCTGATGCTGGTGTCCAGAGAAGGCCCTCATGGGCA	2320						
GTCCGCAAGGCTTCAAGAGCCACGCTCTACTTGTACAGACCTCCAGCCGTACATCGACGATTCGTGGCTCACCTGTCA	2400						

FIG. 5A

8/14

2410	2420	2430	2440	2450	2460	2470	2480
GGAGACGAGCCCGCTGAGGATGCGTGTATCGAGCAGAGCTCTCCCTGATGAGGCGAGTGGCTCTTCGACG	2490	2500	2510	2520	2530	2540	2550
TTCTTCTACGCTTATGTCACACGCGTGGCATCAGGGCAAGTCTCTACGTCCAGTCCAGGGATCCCGCAGGGC	2560	2570	2580	2590	2600	2610	2620
TCCATCCTCTCCACGCTGCTGTGAGCTGTGTACGGCGACATGGAGAACAAAGCTGTTTGGGGATTCGGCGGACGG	2630	2640	2650	2660	2670	2680	2690
GCTGCTCCTGCGTTTGGTGATGATTTCTTGTGGTGACACCTCACCCACCGGAAACCTTCTCAGGACCCCTGG	2700	2710	2720	2730	2740	2750	2760
TCCGAGGTGTCCCTGAGTATGGCTGCGTGGTGAACCTTCCGGAAGACAGTGGTGAACCTTCCCTGTAGAAGACGAGGCCCTG	2770	2780	2790	2800	2810	2820	2830
2840	2850	2860	2870	2880	2890	2900	2910
GGTGGCACGGCTTTGTTAGATGCCGGCCACGGCCATTTCCCTGGTGGCGCTGTCTGGATACCCGGACCCCTGGA	2920	2930	2940	2950	2960	2970	2980
GGTGACAGCGACTACTCCAGCTATGCCCGACCTCCATCAGAGCCAGTCTCACCTTCAACCGCGCTTCAAGGCTGGGA	2990	3000	3010	3020	3030	3040	3050
GGAAACATGCGTCGAAACCTTTTGGGCTTTCGGGCTGAAGTGTACAGCCTGTTTCTGGATTGGCAGGTGAACAGCCTC	3060	3070	3080	3090	3100	3110	3120
CAGACGGTGTGCACCAACATCTACAGATCCTTCTGCTGAGGGCTACAGGTTCACGCATGTGTGCTGCAGCTCCCAT	3130	3140	3150	3160	3170	3180	3190
TCATCAGCAAGTTTGGAGAGACCCCAATTTTCTGCGCGTCACTCTGACACGGCCCTCCCTCTGTCTACTCCATCCTGA	3200	3210	3220	3230	3240	3250	3260
3270	3280	3290	3300	3310	3320	3330	3340
AAGCCAGAACGACGAGGATGTGCTGGGGCCCAAGGGCCCGCGGCCCTCTGCGCTCCGAGGCCGTGCAGTGGCTGTGC	3350	3360	3370	3380	3390	3400	3410
CACCAAGCATTCCTGTCAAGCTGACTCGACACCGTGTACCTACGTGCCACTCCTGGGGTCACTCAGGACAGCCAGAC	3420	3430	3440	3450	3460	3470	3480
GCAGTGAATCGGAAGCTCCCGGGACGACGCTGACTGCCCTGGAGGGCCGAGCAACCCGGCAGTCCCTCAGACTTCA	3490	3500	3510	3520	3530	3540	3550
AGACCATCTTGACTGATGGCCACCCGCCACAGCCAGCCGAGAGCAGACACCAAGGCTGTGTACGCGCGGCTCTAC	3560	3570	3580	3590	3600	3610	3620
GTCCCCAGGGAGGGGGCGGCCACACCCAGGCCCGCACCGCTGGGAGTCTGAGGCCCTGAGTGTGTTTGGCCCGAGGC	3630	3640	3650	3660	3670	3680	3690
3700	3710	3720	3730	3740	3750	3760	3770
CTGCAATGTCGGGTGAAGGCTGAGTGTCCGGCTGAGGCTGAGCGAGTGTCCAGCCCAAGGCTGAGTGTCCAGCACCT	3780	3790	3800	3810	3820	3830	3840
GCCGTCTTCACTTCCCAACAGGTGGCGCTCGGCTCCACCCAGGGCCAGCTTTTCTTCCACAGGAGCCCGGCTTCCACT	3850	3860	3870	3880	3890	3900	3910
CCCCACATAGGAATAGTCCATCCCGAGATTCCGCAATTGTTCACCCCTCGCCCTGCCCTTCTTGCCTTCCACCCCA	3920	3930	3940	3950	3960	3970	3980
TCCAGGTGGAGACCCCTGAGAGGACCCCTGGAGCTCTGGGAAATTGGAGTGACCAAGGTGTGCCCTGTACACAGGCGAG	3990	4000	4010	4020	4030	4040	4050
GACCTGCACCTGGATGGGGTCCCTGTGGGTCAAATTGGGGGAGGTGCTGTGGGAGTAAATACTGAATATATAGATT	4060	4070	4080	4090	4100	4110	4120
4130	4140	4150	4160	4170	4180	4190	4200
TTTTCAGTTTGTGAAAAAATAAAA 4023							

FIG. 5B

9/14

MPRAPRCRAVRSLLRSHYREVLPATFVRRLGPQGWRLVQRGDPAAFRALVAQCLVCVPWDARPPPAAPSF  
 QVSCLEL  
 VARVLQRLCERGAKNVLAFGFALLDGARGGPPPEAFTTSVRSYLPNTVTDALRGSGAWGLLLRRVGGDDVLVH  
 LLARCALFV  
 LVAPSCAYQVCGPPLYQLGAATQARPPPHASGPRRLGCERAWNHSVREAGVPLGLPAPGARRRGGSASRSL  
 PLPKRPRR  
 GAAPERTPVGGSWAHPGRTRGPSDRGFCVSPARPAEEATSEALSGTRHSHPSVGRQHHAGPPSTSRPP  
 RPWDTF  
 CPPVYAEKHFYSSGDKEQLRPSFLLSSLRPSLTGARRLVETIFLGSRPWMPGTFRRLPRLPQRYWQMRPLFL  
 ELLGNH  
 AQCFYGVLLKTHCPLRAAATPAAGVCAREKPOGSVAAPPEEDTDPRRLVQLLRQHSFPWQVYGFVRACLRRL  
 VPPGLWGS  
 RHNERRFLRNTKKEFISLGKHAKLSLQELTWKMSVRGCWLRRSPGVCVPAEAHRLREEILAKFLHWMMSVY  
 VVELLRSE  
 FYVTETTFQKNRLEFFYRKSVMWSKLQSIGIRQHLKRVQLRELSEAEVRQHREARPALLTSRLRFIPKPDGLRPV  
 MDYV  
 GARTFRREKRAERLTSRVKALFVNLNERARRPGLLGASVLGLDDIHRAWRTFVLRVRAQDPPPELYFVKVD  
 VTGAYDTI  
 PQDRLTEVIASIIKPQNTYCVRRYAVVQAAHGHVRKAFKSHVSTLTDLQPYMRQFVAHLQETSFRLDVAVIE  
 QSSSLNE  
 ASSGLFDVFLRFMCHHAVRIRKSYVQCQGIPOGSIILSTLLCSLCYGD MENKLFAGIRRDGLLLRLVDDFLLVT  
 PHLTHA  
 KTFRLTLVRGVPEYGCVVNLKRTVNFPEDEALGGTAFVQMPAHGLFPWCGLLLDTRTLEVSQSYSSYART  
 SIRASLTF  
 NRGFKAGRNMRKLFGLRLKCHSLFLDLQVNSLQTVCTNIYKILLQAYRFHACVLQLPFHQVWKNPTFF  
 LRVISDTA  
 SLCSILKAKNAGMSLGAKGAAGPLPSEAVQWLCHQAFLLKLTTRHRTVTVPLLGSLRTAQTLRSRKLPGTTLT  
 ALEAAN  
 PALPSDFKTILD

FIG. 6

10/14

10	20	30	40	
GTCTCACTCTGTACCCAGGCTGGAGTGTAATGGCACAAT				40
CTCGGCTCACTGCAACCTCTGCCCTCTCGGTTCAAGCAGT				80
TCTCATTCTCAACCTCATGAGTAGCTGGGATTACAGGCG				120
CCCACCACCACGCCTGGTTAATTTTGTATTTTGTAGTAGA				160
GATAGGCTTTTACCATGTTGGCCAGGCTGGTCTCAAATC				200

210	220	230	240	
CTGACCTCAAGTGATCTGCCCGCTTGGCCTCCACAGTG				240
CTGGGATTACAGGTGCAAGCCACCGTGCCCGGCATACCTT				280
GATCTTTTAAATGAAGTCTGAAACATTGCTACCCTTGTC				320
CTGAGCAATAAGACCCCTTAGTGATTTTAGCTCTGGCCAC				360
CCCCCAGCCTGTGTGCTGTTTTCCTGCTGACTTAGTTCTA				400

410	420	430	440	
TCTCAGGCATCTTGACACCCCAAGCTAAGCATTATTA				440
ATATTGTTTTCCGTGTTGAGTGTTTCTGTAGCTTTGCCCC				480
CGCCCTGCTTTTCTCCTTTGTTCCCGTCTGTCTTCTGT				520
CTCAGGCCCCGCGTCTGGGGTCCCTTCCTTGCTCCTTGC				560
GTGGTCTTCTGTCTTGTTATTGCTGGTAAACCCAGCTT				600

610	620	630	640	
TACCTGTGCTGGCCTCCATGGCATCTAGCGACGTCCGGGG				640
ACCTCTGCTTATGATGCACAGATGAAGATGTGGAGACTCA				680
CGAGGAGGGCGGTATCTTGGCCCGTGAGTGCTGGAGCA				720
CCACGTGGCCAGCGTTCCCTTAGCCAGTGAGTGACAGCAAC				760
GTCCGCTCGGCCTGGGTTTCAGCCTGAAAACCCAGGCAN				800

810	820	830	840	
GTCGGGGTCTGGTGGCTCCGCGGTGTCGAGTTTGAAATCG				840
CGCAAACCTGCGGTGTGGCGCCAGCTCTGACGGTGCTGCC				880
TGGCGGGGGAGNGTCTGCTTCCTCCCTTCTGCTTGGGAAC				920
CAGGACAAAGGATGAGGCTCCGAGCCGTTGTGCCCCAACA				960
GGAGCATGACGGGTTGGCTGTGTTCCGGCCGAGAGCACC				1000

1010	1020	1030	1040	
GTCTGCGTGAGGAGATCCTGGCCAAGTTCCTGCACTGGCT				1040
GATGAGTGTGTACGTCGTCGAGCTGCTCAGGTCTTTCTTT				1080
TATGTCACGGAGACCACGTTTCAAAGAAGAGGCTCTTTT				1120
TCTACCGGAAGAGTGTCTGGAGCAAGTTGCAAAGCATGG				1160
AATCAGACAGCACTTGAAGAGGGTGAGCTGCGGGAGCTG				1200

FIG. 7A

11/14

1210	1220	1230	1240
TCGGAAGCAGAGGTCAGGCAGCATCGGGAAGCCAGGCCCG			1240
CCCTGCTGACGTCCAGACTCCGCTTCATCCCCAAGCCTGA			1280
CGGGCTGCGGCCGATTGTGAACATGGACTACGTCGTGGGA			1320
GCCAGAACGTTCCGCAGAGAAAAGAGGGCCGAGCGTCTCA			1360
CCTCGAGGGTGAAGGCACTGTTTCAGCGTGCTCAACTACGA			1400
1410	1420	1430	1440
GCGGGCGCGCGGCCCGGCCCTCCTGGGCGCCTCTGTGCTG			1440
GGCCTGGACGATATCCACAGGGCCTGGCGCACCTTCGTGC			1480
TGCGTGTGCGGGCCAGGACCCGCCGCTGAGCTGTACTT			1520
TGTCAAGGTGGATGTGACGGGCGGTACGACACCATCCCC			1560
CAGGACAGGCTCACGGAGGTCATCGCCAGCATCATCAAAC			1600
1610	1620	1630	1640
CCCAGAACACGTACTGCGTGCGTCCGTATGCCGTGGTCCA			1640
GAAGGCCGCCCATGGGCACGTCCGCAAGGCCCTTCAAGAGC			1680
CACGTCTACGTCCAGTGCCAGGGGATCCCGCAGGGCTCC			1720
ATCCTCTCCACGTGCTCTGCAGCCTGTGCTACGGCGACA			1760
TGGAGAACAAAGCTGTTTGCGGGGATTTCGGCGGGACGGGCT			1800
1810	1820	1830	1840
GCTCCTGCGTTTGGTGGATGATTTCTTGTGGTGACACCT			1840
CACCTCACCCACGCGAAACCTTCCTCAGGACCCCTGGTCC			1880
GAGGTGTCCCTGAGTATGGCTGCGTGGTGAACCTGCGGAA			1920
GACAGTGGTGAACCTCCCTGTAGAAGACGAGGCCCTGGGT			1960
GGCACGGCTTTTGTTCAGATGCCGGCCACGGCCTATTC			2000
2010	2020	2030	2040
CCTGGTGCGGCCCTGCTGCTGGATAACCGGACCCCTGGAGGT			2040
GCAGAGCGACTACTCCAGCTATGCCCGGACCTCCATCAGA			2080
GCCAGTCTCACCTTCAACCGCGGCTTCAAGGCTGGGAGGA			2120
ACATGCGTCGCAAACTCTTTGGGGTCTTGCGGCTGAAGTG			2160
TCACAGCCTGTTTCTGGATTTCAGGTGAACAGCCTCCAG			2200
2210	2220	2230	2240
ACGGTGTGCACCAACATCTACAAGATCCTCCTGCTGCAGG			2240
CGTACAGGTTTTCACGCATGTGTGCTGCAGCTCCCATTTCA			2280
TCAGCAAGTTTGAAGAACCCACATTTTCTGCGCGTC			2320
ATCTCTGACACGGCCTCCCTCTGCTACTCCATCCTGAAAG			2360
CCAAGAACGCAGCCGAAGAAAACATTTCTGTGCTGACTCC			2400
2410	2420	2430	2440
TGCGGTGCTTGGGTGCGGACAGCCAGAGATGGAGCCACCC			2440
CGCAGACCGTCGGGTGTGGGCAGCTTTCGGTGTCTCCTG			2480
GGAGGGGAGCTGGGCTGGGCCTGTGACTCCTCAGCCTCTG			2520
TTTTCCCCCAGGGATGTGCTGGGGGCCAAGGGCGCCGCC			2560
GGCCCTCTGCCCTCCGAGGCCGTGCAGTGGCTGTGCCACC			2600

FIG. 7B

12/14


2610	2620	2630	2640
AAGCATTCCTGCTCAAGCTGACTCGACACCGTGTCACCTA			2640
CGTGCCACTCCTGGGGTCACTCAGGACAGCCCAGACGCAG			2680
CTGAGTCGGAAAGCTCCCGGGGACGACGCTGACTGCCCTGG			2720
AGGCCGCAGCCAACCCGGCACTGCCCTCAGACTTCAAGAC			2760
CATCCTGGACTGATGGCCACCCGCCACAGCCAGGCCGAG			2800

2810	2820	2830	2840
AGCAGACACCAGCAGCCCTGTACGCCGGGCTCTACGTCC			2840
CAGGGAGGGAGGGGCGGCCACACCCAGGCCCGCACCGCT			2880
GGGAGTCTGAGCCTGAGTGAGTGTTTGCCGAGGCCTGC			2920
ATGTCCGGCTGAAGGCTGAGTGTCGGCTGAGGCCTGAGC			2960
GAGTGTCAGCCAAGGGCTGAGTGTCAGCACACCTGCCG			3000

3010	3020	3030	3040
TCTTCACTTCCCCACAGGCTGGCGCTCGGCTCCACCCAG			3040
GGCCAGCTTTTCCTCACCAGGAGCCCGGCTTCACTCCCC			3080
ACATAGGAATAGTCCATCCCCAGATTCGCCATTGTTACC			3120
CCTCGCCCTGCCCTCCTTTGCCTTCCACCCCAACCATCCA			3160
GGTGGAGACCCCTGAGAAGGACCCCTGGGAGCTCTGGGAATT			3200

3210	3220	3230	3240
TGGAGTGACCAAAGGTGTGCCCTGTACACAGGCGAGGACC			3240
CTGCACCTGGATGGGGGTCCCTGTGGGTCAAATTGGGGGG			3280
AGGTGCTGTGGGAGTAAAATACTGAATATATGAGTTTTTC			3320
AGTTTTGAAAAAAAAAAAAAGGAATTC			3346

FIG. 7C





14/14

FIG. 9A

FIG. 9B




FIG. 9C

FIG. 9D